



Author(s)	Shattuck, Nita Lewis; Matsangas, Panagiotis
Title	Work and rest patterns and psychomotor vigilance performance of crewmembers of the USS Jason Dunham: a comparison of the 3/9 and 6/6 watchstanding schedules
Publisher	
Issue Date	2014 12
URL	<a href="http://hdl.handle.net/10945/44348">http://hdl.handle.net/10945/44348</a>

This document was downloaded on February 09, 2015 at 12:49:50



<http://www.nps.edu/library>

Calhoun is a project of the Dudley Knox Library at NPS, furthering the precepts and goals of open government and government transparency. All information contained herein has been approved for release by the NPS Public Affairs Officer.

**Dudley Knox Library / Naval Postgraduate School  
411 Dyer Road / 1 University Circle  
Monterey, California USA 93943**



<http://www.nps.edu/>



# NAVAL POSTGRADUATE SCHOOL

MONTEREY, CALIFORNIA

**WORK AND REST PATTERNS AND PSYCHOMOTOR  
VIGILANCE PERFORMANCE OF CREWMEMBERS OF THE  
USS JASON DUNHAM: A COMPARISON OF THE 3/9 AND 6/6  
WATCHSTANDING SCHEDULES**

by

Nita Lewis Shattuck, Ph.D. and Panagiotis Matsangas, Ph.D.

December 2014

**Approved for public release; distribution is unlimited**

Prepared for: United States Navy, N171, 21<sup>st</sup> Century Sailor Office and the Naval Medical Research Center, Advanced Medical Development Office, 503 Robert Ave., Bldg. 500, Silver Spring, MD 20910

THIS PAGE INTENTIONALLY LEFT BLANK

<b>REPORT DOCUMENTATION PAGE</b>			<i>Form Approved</i> OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
<b>1. REPORT DATE (DD-MM-YYYY)</b> 31-12-2014		<b>2. REPORT TYPE</b> Technical Report		<b>3. DATES COVERED (From-To)</b> October 2013 – September 2014	
<b>4. TITLE AND SUBTITLE</b> WORK AND REST PATTERNS AND PSYCHOMOTOR VIGILANCE PERFORMANCE OF CREWMEMBERS OF THE USS JASON DUNHAM: A COMPARISON OF THE 3/9 AND 6/6 WATCHSTANDING SCHEDULES			<b>5a. CONTRACT NUMBER</b>		
			<b>5b. GRANT NUMBER</b>		
			<b>5c. PROGRAM ELEMENT NUMBER</b>		
<b>6. AUTHOR(S)</b> Nita Lewis Shattuck, Ph.D. and Panagiotis Matsangas, Ph.D.			<b>5d. PROJECT NUMBER</b>		
			<b>5e. TASK NUMBER</b>		
			<b>5f. WORK UNIT NUMBER</b>		
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) AND ADDRESS(ES)</b> Operations Research Department Naval Postgraduate School Monterey, CA 93943			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b> NPS-OR-14-004		
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> United States Navy, N171, 21 <sup>st</sup> Century Sailor Office, Naval Medical Research Center Advanced Medical Development Office, 503 Robert Ave Bldg. 500, Silver Spring, MD 20910			<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>		
			<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>		
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for public release; distribution is unlimited.					
<b>13. SUPPLEMENTARY NOTES</b> The views expressed in this report are those of the author(s) and do not reflect the official policy or position of the Department of Defense or the U.S. Government.					
<b>14. ABSTRACT</b> This study compares the patterns of crew rest and sleep, psychomotor vigilance performance, and work demands/rest opportunities afforded by two different schedules, the 3-hour on/9-hour off ("3/9") and the 6-hour on/6-hour off ("6/6") watchstanding schedules. The study was conducted aboard the USS Jason Dunham, a U.S. Navy destroyer operating in the vicinity of the Persian Gulf during the months of November and December 2012. Of the 122 participants in the overall study, 52 were shift workers using either the 3/9 (n=41) or the 6/6 (n=11) schedules. These 52 individuals are the focus of the current analysis. Although sleep deprivation was evident in both watch schedules, results show that crewmembers on the 3/9 received more sleep than their peers on the 6/6, with 6.46 ±0.77 hours versus 5.89 ±0.87 hours, respectively. The 3/9 schedule, compared to the 6/6, was also better in terms of the distribution of sleep episodes across the day. Specifically, crewmembers on the 3/9 received more sleep during nighttime hours, whereas crewmembers on the 6/6 had to sleep during the day to compensate for their lack of sleep during nighttime hours. In terms of work demands, crewmembers on the 6/6 schedule have considerably long workdays, with, on average, 15 hours on duty, which corresponds to approximately 30% more time on duty than allocated in the Navy Standard Work Week criterion (on average, 105 hours compared to 81 hours weekly). The two schedules differed significantly in the variability of psychomotor vigilance performance; specifically, crewmembers on the 6/6 schedule had larger variability than those on the 3/9 in 11 of the 13 Psychomotor Vigilance Task (PVT) metrics analyzed (p<0.05). The average value of the PVT scores was better on the 3/9 compared to the 6/6, but not at statistically significant levels. The findings of this study show that the 3/9 is better than the 6/6 in affording rest and sleep opportunities, sleep hygiene, fatigue levels, psychomotor vigilance performance, work demands, and acceptance from the participants.					
<b>15. SUBJECT TERMS</b> Watch schedules, sleep, psychomotor vigilance performance					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b> UU	<b>18. NUMBER OF PAGES</b> 61	<b>19a. NAME OF RESPONSIBLE PERSON</b> Nita Lewis Shattuck
<b>a. REPORT</b> Unclassified	<b>b. ABSTRACT</b> Unclassified	<b>c. THIS PAGE</b> Unclassified			<b>19b. TELEPHONE NUMBER (include area code)</b> (831) 656-2281

Standard Form 298 (Rev. 8-98)  
Prescribed by ANSI Std. Z39.18

THIS PAGE INTENTIONALLY LEFT BLANK

**NAVAL POSTGRADUATE SCHOOL  
Monterey, California 93943-5000**

Ronald A. Route  
President

Douglas A. Hensler  
Provost

The report entitled “*Work and Rest Patterns and Psychomotor Vigilance Performance of Crewmembers of the USS Jason Dunham: A Comparison of the 3/9 and 6/6 Watchstanding Schedules*” was prepared for and funded by United States Navy, N171, 21<sup>st</sup> Century Sailor Office and the U.S. Navy Bureau of Medicine Advanced Medical Devices.

**Further distribution of all or part of this report is authorized.**

**This report was prepared by:**

Nita Lewis Shattuck, Ph.D.  
Associate Professor of  
Operations Research

Panagiotis Matsangas, Ph.D.  
National Research Council  
Postdoctoral Associate

**Reviewed by:**

Robert F. Dell, Chairman  
Chairman  
Department of Operations Research

Johannes O. Royset  
Associate Chairman for Research  
Department of Operations Research

**Released by:**

Jeffrey D. Paduan  
Dean of Research

THIS PAGE INTENTIONALLY LEFT BLANK

## ABSTRACT

This study compares the patterns of crew rest and sleep, psychomotor vigilance performance, and work demands/rest opportunities afforded by two different schedules, the 3-hour on/9-hour off (“3/9”) and the 6-hour on/6-hour off (“6/6”) watchstanding schedules. The study was conducted aboard the USS Jason Dunham, a U.S. Navy destroyer operating in the vicinity of the Persian Gulf during the months of November and December 2012. Of the 122 participants in the overall study, 52 were shift workers using either the 3/9 (n=41) or the 6/6 (n=11) schedules. These 52 individuals are the focus of the current analysis. Although sleep deprivation was evident in both watch schedules, results show that crewmembers on the 3/9 received more sleep than their peers on the 6/6, with  $6.46 \pm 0.77$  hours versus  $5.89 \pm 0.87$  hours, respectively. The 3/9 schedule, compared to the 6/6, was also better in terms of the distribution of sleep episodes across the day. Specifically, crewmembers on the 3/9 received more sleep during nighttime hours, whereas crewmembers on the 6/6 had to sleep during the day to compensate for their lack of sleep during nighttime hours. In terms of work demands, crewmembers on the 6/6 schedule have considerably long workdays, with, on average, 15 hours on duty, which corresponds to approximately 30% more time on duty than allocated in the Navy Standard Work Week (NSWW) criterion (on average, 105 hours compared to 81 hours weekly).

The two schedules differed significantly in the variability of psychomotor vigilance performance; specifically, crewmembers on the 6/6 schedule had larger variability than those on the 3/9 in 11 of the 13 Psychomotor Vigilance Task (PVT) metrics analyzed ( $p < 0.05$ ). The average value of the PVT scores was better on the 3/9 compared to the 6/6, but not at statistically significant levels. The findings of this study show that the 3/9 is better than the 6/6 in affording rest and sleep opportunities, sleep hygiene, fatigue levels, psychomotor vigilance performance, work demands, and acceptance from the participants.



THIS PAGE INTENTIONALLY LEFT BLANK

# TABLE OF CONTENTS

<b>I.</b>	<b>INTRODUCTION.....</b>	<b>1</b>
<b>A.</b>	<b>SHIFT WORK IN THE NAVAL ENVIRONMENT .....</b>	<b>4</b>
<b>B.</b>	<b>NAVY STANDARD WORK WEEK (NSWW) MODEL .....</b>	<b>6</b>
<b>C.</b>	<b>SCOPE .....</b>	<b>7</b>
<b>II.</b>	<b>METHODOLOGY .....</b>	<b>9</b>
<b>A.</b>	<b>EXPERIMENTAL DESIGN .....</b>	<b>9</b>
<b>B.</b>	<b>PARTICIPANTS.....</b>	<b>9</b>
<b>C.</b>	<b>EQUIPMENT AND INSTRUMENTS.....</b>	<b>9</b>
<b>1.</b>	<b>Actiwatches.....</b>	<b>9</b>
<b>2.</b>	<b>Epworth Sleepiness Scale (ESS) .....</b>	<b>10</b>
<b>3.</b>	<b>Morningness-Eveningness (ME) Preference .....</b>	<b>10</b>
<b>4.</b>	<b>Pittsburgh Sleep Quality Index (PSQI) .....</b>	<b>10</b>
<b>5.</b>	<b>Psychomotor Vigilance Test (PVT) .....</b>	<b>11</b>
<b>6.</b>	<b>Activity Logs.....</b>	<b>11</b>
<b>D.</b>	<b>PROCEDURES .....</b>	<b>11</b>
<b>E.</b>	<b>VARIABLES .....</b>	<b>12</b>
<b>F.</b>	<b>ANALYSIS .....</b>	<b>12</b>
<b>1.</b>	<b>Actigraphic Data Cleaning and Reduction Procedures .....</b>	<b>12</b>
<b>2.</b>	<b>PVT Data Cleaning and Reduction Procedures.....</b>	<b>14</b>
<b>3.</b>	<b>Sleep Log Data Cleaning and Reduction Procedures for Workload Analysis .....</b>	<b>14</b>
<b>4.</b>	<b>Analysis Roadmap .....</b>	<b>15</b>
<b>III.</b>	<b>RESULTS .....</b>	<b>17</b>
<b>A.</b>	<b>DEMOGRAPHICS .....</b>	<b>17</b>
<b>B.</b>	<b>DESCRIPTIVE RESULTS FOR THE ENTIRE DATA SET .....</b>	<b>18</b>
<b>C.</b>	<b>PRIMARY ANALYSIS .....</b>	<b>19</b>
<b>1.</b>	<b>Comparison Between Watch Sections.....</b>	<b>21</b>
<b>2.</b>	<b>Psychomotor Vigilance Performance.....</b>	<b>24</b>
<b>3.</b>	<b>Posttest Questionnaires .....</b>	<b>26</b>
<b>4.</b>	<b>Work Patterns .....</b>	<b>27</b>
<b>IV.</b>	<b>DISCUSSION .....</b>	<b>33</b>
<b>A.</b>	<b>SOME THOUGHTS ON WORK DEMANDS AND REST AFFORDABILITY .....</b>	<b>36</b>
<b>B.</b>	<b>CAVEATS .....</b>	<b>37</b>
	<b>APPENDIX. PARTICIPANT INFORMATION.....</b>	<b>39</b>
	<b>LIST OF REFERENCES.....</b>	<b>41</b>
	<b>INITIAL DISTRIBUTION LIST .....</b>	<b>47</b>

THIS PAGE INTENTIONALLY LEFT BLANK

## LIST OF FIGURES

Figure 1.	Effects of shift work on individuals and organizations. ....	4
Figure 2.	Daily rest amount in hours. Numbers on vertical bars represent the corresponding number of participants. ....	18
Figure 3.	Daily sleep amount in hours. Numbers on vertical bars represent the corresponding number of participants. ....	19
Figure 4.	Daily sleep and number of sleep episodes per day by watch schedule.....	20
Figure 5.	Epworth Sleepiness Scale scores by watch schedule.....	21
Figure 6.	Daily sleep and the number of sleep episodes per day in the 3/9 watch sections.....	22
Figure 7.	Daily rest and sleep, by watch schedule and shift. ....	24
Figure 8.	Differences in standard deviations in PVT metrics between the 3/9 and 6/6 watch schedules. ....	26
Figure 9.	Subjective evaluations of watch schedules. ....	27
Figure 10.	Work and sleep patterns in the 00:00 – 03:00 and 12:00 – 15:00 section of the 3/9 watch schedule. ....	29
Figure 11.	Work and sleep patterns in the 03:00 – 06:00 and 15:00 – 18:00 section of the 3/9 watch schedule. ....	29
Figure 12.	Work and sleep patterns in the 06:00 – 09:00 and 18:00 – 21:00 section of the 3/9 watch schedule. ....	30
Figure 13.	Work and sleep patterns in the 09:00 – 12:00 and 21:00 – 23:59 section of the 3/9 watch schedule. ....	30
Figure 14.	Work and sleep patterns in the 00:00 – 06:00 and 12:00 – 18:00 section of the 6/6 watch schedule. ....	31
Figure 15.	Work and sleep patterns in the 06:00 – 12:00 and 18:00 – 23:59 section of the 6/6 watch schedule. ....	31
Figure 16.	Polar diagrams of typical 24-hour days in the 3/9 (left diagram) and 6/6 watch schedules (right diagram). ....	32

THIS PAGE INTENTIONALLY LEFT BLANK

## LIST OF TABLES

Table 1.	Demographics. ....	17
Table 2.	Daily rest and sleep amount by participant. ....	18
Table 3.	Daily rest/sleep amount in hours and number of sleep episodes per day by watch schedule. ....	20
Table 4.	Daily rest/sleep amount in hours by watch shift. ....	23
Table 5.	Comparison of PVT metrics between the 3/9 and 6/6 watch schedules. ....	25
Table 6.	Work patterns by watch schedule. ....	27
Table 7.	Comparison ranks between watch schedules. ....	35
Table 8.	Participants by watch schedule, watch section, and actiwatch type. ....	39

THIS PAGE INTENTIONALLY LEFT BLANK

## I. INTRODUCTION

Even though shift work is an integral part of modern societies, it remains an abnormal behavior for humans. Many human physiological functions are controlled by the circadian clock; for example, sleep and its associated functions are promoted during the biological night, whereas wakefulness is more common during the biological day (Dijk & Czeisler, 1995; Dijk & Edgar, 1999; Drake & Wright, 2011). Research findings suggest that shift work, that is, working hours other than the typical daytime work hours of 8:00 am to 5:00 pm, affects physiological function and performance through circadian desynchrony. This disruption of the internal circadian rhythm (Colquhoun, Blake, & Edwards, 1969b) is accompanied by an alteration of melatonin production (Burch, Yost, Johnson, & Allen, 2005) and disturbance of the sleep-wakefulness cycle (Åkerstedt, 2003). The brain undergoes long-term changes when subjected to chronic sleep restriction (Belenky et al., 2003) altering neural function in a manner that precludes rapid recovery to baseline levels of alertness and performance, even after sleep durations return to baseline levels. The time needed to adjust to a nocturnal rhythm is at least a week (Monk, 1986), although 12 days or more have also been reported (Colquhoun, Blake, & Edwards, 1969a; Hockey, 1983). Workers often experience sleep disturbance and sleepiness after months or years of shift work (Drake & Wright, 2011). Research findings suggest that prior exposure to shift work is related to sleep problems during retirement (Monk et al., 2013). Irrespective of the shift system, night and early-morning shifts are associated with short sleep and increases in sleepiness (Sallinen & Kecklund, 2010). Compared to non-watchstanders or day workers, shift work results in more sleep deprivation, shorter sleep episodes, greater sleep fragmentation, and increased levels of fatigue (Arendt, Middleton, Williams, Francis, & Luke, 2006).

Long duration shifts—that is, shifts greater than 12 hours in length—lead to increased sleepiness due to reduced sleep opportunities afforded to shift workers (Åkerstedt & Wright, 2009). Short sleep and increased sleepiness are also evident in individuals working long shifts (i.e., >16 hours) and extended weekly working hours (i.e., >55 hours) (Sallinen & Kecklund, 2010).



In general, researchers agree that forward-rotating shifts are better for alertness and sleep (Hockey, 1983; Sallinen & Kecklund, 2010; Viitasalo, Kuosma, Laitinen, & Härmä, 2008), even though some studies do not fully support the argument that rotating clockwise is better than rotating counterclockwise (Cruz, Boquet, Detwiler, & Nesthus, 2003). Many studies show that rotating watch systems have a detrimental impact on the sleep/wake cycle (Colquhoun & Folkard, 1985; Goh, Tong, Lim, Low, & Lee, 2000; Hakola & Härmä, 2001); however, there are conflicting results regarding the utility of fixed-shift systems compared to rotating systems (Åkerstedt, 2003; Folkard, 1992; Wedderburn, 1992; Wilkinson, 1992). Supporters of fixed systems suggest that, given adequate time, the human circadian pacemaker will adjust to night shift work, whereas other researchers argue that humans can never fully adjust their sleep/wake patterns to night shift work because of social or other factors (Colquhoun & Folkard, 1985; Cruz et al., 2003; Monk, 1990). Studies show that shift work can lead to considerable performance degradation (as measured by sustained attention, vigilance, and simulated driving tasks) caused by sleepiness at levels equivalent to 0.04 to 0.05 g% blood alcohol concentration (Arnedt, Owens, Crouch, Stahl, & Carskadon, 2005).

Shift work is also associated with elevated levels of physiological fatigue in manual tasks involving the upper extremities (Rosa, Bonnet, & Cole, 1998). Specifically, longer shifts (12 hours compared to 8 hours) and night shifts lead to increased fatigue. Shiftwork, however, not only leads to circadian desynchrony, sleep deprivation, and performance impairment, but it also has considerable negative social consequences on balancing work and home life. Scientific evidence suggests that shift work has a negative influence on children's well-being and on marital satisfaction (Albertsen, Rafnsdóttir, Grimsmo, Tómasson, & Kauppinen, 2008). Shift work is also associated with weight gain and various morbidities. A study assessed the health effects of implementing a 12-hour shift in place of the traditional 8-hour shift in factory workers in Japan (Yamada, Tachibana, & Kuriyama, 1988). Compared to the workers on the 8-hour shift, the workers changing to a 12-hour shift had significant increases in psychological symptoms related to fatigue and experienced, on average, a weight gain of one kilogram. Research findings also suggest that shift workers are more prone to developing other disorders, e.g., obesity, gastrointestinal disorders, cardiovascular heart disease, compromised

pregnancy outcome, breast cancer, prostate cancer, metabolic syndrome, and diabetes (Drake & Wright, 2011; Folkard & Tucker, 2003; Harrington, 2001; Knutsson, 2003; Wang, Armstrong, Cairns, Key, & Travis, 2011). The International Agency for Research on Cancer has classified “shift work that involves circadian disruption” as a probable human carcinogen (Stevens et al., 2011, p. 764).

Shift work can have a significant impact on organizational risk and safety rates. Epidemiological studies have shown that the relative risk of workplace accidents increases by approximately 15% during afternoon shifts and by 30% for night shifts compared to morning shifts (Folkard, Lombardi, & Spencer, 2006). Long work hours and shift work increase the risk of workplace accidents (Åkerstedt & Wright, 2009), with accident rates increasing between 50% and 100% (Wagstaff & Sigstad Lie, 2011). Work periods of more than 8 hours carry an increased risk of accidents that cumulates, so that the increased risk of accidents at around 12 hours is twice the risk at 8 hours. The authors noted, however, that “pure” night work may bring some protection against the detrimental effects of shift work (Wagstaff & Sigstad Lie, 2011).

Knutsson (1989, 2003) proposed a conceptual model to explain mechanisms of disease in shift workers by integrating relevant research findings. Although the initial model focused only on disease, it can be extended to include the effect on human performance as well as on organizational health and resilience (Rutenfranz, Colquhoun, Knauth, & Ghata, 1977). Figure 1 demonstrates a revision to the initial model.




Figure 1. Effects of shift work on individuals and organizations.

Overall, optimizing a shift schedule for a specific work environment is not an easy task; a large number of parameters must be considered. Given the complexity in assessing the utility of a watch system, Miller (2006) proposed nine “principles” of shift-work scheduling, in the sense that these “principles” describe essential qualities of shift systems. These nine principle were classified into three groups: circadian stability, principle of chronohygiene (short shift length, minimum number of consecutive night shifts, recovery after each night shift, maximum number of free days on weekends, at least 104 days off per year), and principles of satisfaction (equity among shift workers for types of work dates and free days, predictability of specific work and free days, and quality of time off).

#### A. SHIFT WORK IN THE NAVAL ENVIRONMENT

The naval environment is characterized by sleep problems, sleep deprivation, and increased levels of fatigue (Howarth, Pratt, & Tepas, 1999). Shift work is considered one of the major factors leading to sleep problems. It is not surprising, therefore, that

optimizing shift work practices is a matter of concern for many navies and has long been a topic of investigation. While at sea, watches must be manned around the clock; consequently, watches are either fixed (i.e., crewmembers work the same time each day), rapidly rotating (i.e., crewmembers work different times every day), or irregular. The watch system to be used depends on the organizational culture, the prior experience of the command leadership, and the number of crewmembers available to stand watch. This final factor is a critical consideration on ships with limited crew size. Studies on naval vessels have shown that watch schedules traditionally used at sea lead to sleep deprivation, sleep fragmentation, suboptimal performance, and worrisome levels of alertness (Paul, Ebisuzaki, McHarg, Hursh, & Miller, 2012; Rutenfranz et al., 1988).

In the U.S. Navy, it is the responsibility of the officer of the watch to ensure that watchstanders are able to stand an effective watch (Department of the Navy, 2012). Given the availability of personnel, the watch itself, and other daily activities, a number of fixed and rotating watch systems are used like the 4-hour on/8-hour off, 6-hour on/6-hour off (“6/6”), the 12-hour on/12-hour off (“12/12”), the 6-hour on/18-hour off (“6/18”), or the 3-hour on/9-hour off (“3/9”). Some of these schedules result in days that are other than 24 hours in length. For example the 5-hour on/10-hour off (“5/10”) is either 15 or 30 hours in length, and the 5-hour on/15-hour off (“5/15”) results in a 20-hour day. Following a 1969 Naval Postgraduate School master’s thesis by Stolgitis (1969), U.S. Navy submarine crews adopted a 6-hour on/12-hour off (“6/12”), three-section watchstanding schedule that results in an 18-hour day.

Several studies have modeled the effects of specific watch schedules by using the Fatigue Avoidance Scheduling Tool (FAST) (Hursh et al., 2004; Kronauer & Stone, 2004). A study in the Royal Canadian Navy (RCN) compared the predicted psychomotor performance of crewmembers on four watch schedules: the 3-section 8-hour on/16-hour off, the 3-section 4-hour on/8-hour off, and the 2-section 8-hour on/4-hour off/4-hour on/8-hour off (“8/4/4/8”), and the 2-section 6/6 schedule used by RCN submarines (Paul, Hursh, & Miller, 2010). The results showed that the 8-hour on/16-hour off, 4-hour on/8-hour off, and the 8/4/4/8 resulted in better performance as compared to the 6/6.

A number of studies have been conducted at the Naval Postgraduate School to assess the effect of shift work on sleep hygiene and the performance of U.S. Navy crews.

Osborn (2004) compared the 6-hour on/12-hour off used in submarines with a 6-hour watch schedule rotating every three days. Although his results showed that the schedule under investigation did not allow the crew to get more sleep, he noted that further research should identify a watch schedule that would provide more sleep, while still accommodating the constraints of submarine operations. Yokeley (2012), in a within-subjects study of sleep and performance of crewmembers onboard a U.S. Navy destroyer, compared performance of crewmembers while working the 3/9 compared to their performance while working the 5/15. The results, based on actigraphic data, showed that crewmembers on the 3/9 ( $6.11 \pm 0.852$ ) got 0.14 hours more sleep ( $p=0.062$ ) on a daily basis than while on the 5/15 ( $5.56 \pm 0.788$ ). In terms of variability, psychomotor vigilance performance of crewmembers in the 3/9 was more consistent (decreased standard deviation of reaction time) compared to the 5/15.

## **B. NAVY STANDARD WORK WEEK (NSWW) MODEL**

The NSWW model is part of the United States Navy Total Force Manpower Policies Procedures and is described in Naval Operations (OPNAV) Instruction 1000.16K (Department of the Navy, 2007). The NSWW represents a standardized version of one week of work performed by a single enlisted Sailor while at sea, and is used to calculate manning levels, which are a theoretical reflection of the minimum manpower resources necessary to accomplish the ship's mission. The workweek for sea duty is a guideline for sustained personnel utilization based on the operational requirements under projected wartime conditions with units in Condition III steaming, as described in OPNAV Instruction 1000.16K, page C-1 (Department of the Navy, 2007). Although not prescriptive, the instruction notes that extending work hours on a routine basis could adversely affect morale, retention, safety, etc., and, as a policy, habitually extending work hours should be avoided (Department of the Navy, 2007).

The NSWW provides guidelines for the time available per person to accomplish the required workload, including watches expressed in average hours per week. The week is divided into two categories, On Duty (or Available) time (81 hours) and Nonavailable time (87 hours). On Duty time refers to the time periods where personnel are occupied by their required duties: watchstanding (56 hours), work (14 hours),

training (7 hours), and service diversion (4 hours). Training contributes to combat readiness and includes activities such as general drills and engineering casualty damage control. Service diversion includes quarters, inspections, sick call, and administrative requirements. Productive Work time (70 hours) includes watchstanding and work. Nonavailable time is comprised of all personal time that is allotted to sleep (56 hours), messing (14 hours), personal needs (14 hours), and free time (3 hours).

Multiple studies conducted at the Naval Postgraduate School have shown that crewmembers work longer hours and sleep less than what is allocated in the NSW model, suffering from significant sleep deprivation (Green, 2009; Haynes, 2007; Mason, 2009). Specifically, Haynes (2007) found that crewmembers worked approximately 14 hours per day, with 85% of them exceeding the 81 hours allotted by the NSW, whereas Green (2009) identified that sailors worked 12.5 hours per day, with 61% of her participants exceeding the NSW model. Mason (2009) found that Senior Chief Petty Officers and Chief Petty Officers averaged 6.26 hours of sleep, while senior officers (Lieutenant Commanders and above) slept approximately 6.4 hours per day.

### **C. SCOPE**

This report compares the 3-hour on/9-hour off and the 6-hour on/6-hour off watch schedules in terms of crew rest and sleep patterns, psychomotor vigilance performance, and work demands/rest opportunities. This work is part of a multiyear effort at the Naval Postgraduate School designed to systematically and empirically assess the wide range of watch schedules used on U.S. Navy ships to provide insight and guidance for future naval operations.

THIS PAGE INTENTIONALLY LEFT BLANK

## **II. METHODOLOGY**

### **A. EXPERIMENTAL DESIGN**

This study used a quasi-experimental approach in which volunteers from various ship departments were recruited to participate in a study of sleep and performance on a U.S. Navy warship.

### **B. PARTICIPANTS**

Participants were volunteers from the USS JASON DUNHAM (DDG 109), an Arleigh Burke class destroyer, Flight IIA (9,300 tons). Initially, 122 crewmembers volunteered to participate in this study. Crewmembers had been working the same schedule for several weeks before the data collection commenced.

### **C. EQUIPMENT AND INSTRUMENTS**

#### **1. Actiwatches**

Two types of actigraphs were used, the Ambulatory Monitoring, Inc. (AMI) Motionlogger Watch and the Philips Respironics (PR) Spectrum actiwatch. Data for both devices were collected in 1-minute epochs. AMI data (collected in the Zero-Crossing Mode) were scored using Action W version 2.7.2155 software. The Cole-Kripke algorithm, with rescoring rules, was used. Sleep statistics criteria for long sleep and long wake episodes was five minutes. The sleep latency criterion was no more than 1-minute wake in 20 minutes period (all values are default for this software). PR data were scored using Actiware software version 6.0.0 (Phillips Respironics, Bend, Oregon). The medium sensitivity threshold (40 counts per epoch) was used, with 10 immobile minutes as the criterion for sleep onset and sleep end (all values are default for this software). Previous research has shown that AMI data analyzed with Cole-Kripke and PR data analyzed with medium sensitivity parameters assess total sleep time for an approximately 8-hour night sleep episode with three minutes precision (average results compared to polysomnography derived 436 minutes of sleep) (Meltzer, Walsh, Traylor, & Westin, 2012). A comparison failed to identify any significant differences between AMI and PR



actiwatches in 3/9 daily rest, daily total sleep time, occurring within watch periods (Wilcoxon Rank Sum test, for all differences  $p > 0.180$ ).

## **2. Epworth Sleepiness Scale (ESS)**

The ESS is a widely used instrument to assess average daytime sleepiness (Johns, 1991). Using a 4-point Likert scale, the individual indicates the chance of dozing off or falling asleep in eight different everyday situations. Scoring of the answers is 0 to 3, with 0 being “would never doze,” 1 is “slight chance of dozing,” 2 is “moderate chance of dozing,” and 3 denotes “high chance of dozing.” Participants are instructed to rate each question according to his/her usual way of life in recent times. Responses are summed to obtain a total score. A sum of 10 or more reflects above-average daytime sleepiness and the need for further evaluation (Johns, 1992). The questionnaire has a high level of internal consistency, as measured by Cronbach’s alpha, ranging from 0.73 to 0.88 (Johns, 1992).

## **3. Morningness-Eveningness (ME) Preference**

The Morningness-Eveningness Scale (Horne & Östberg, 1976) was used to assess participants’ chronotype; an individual human attribute related to whether there is a preference for waking earlier or later in the day. The scale includes 19 multiple-choice questions. Scores range from 16 to 86, with scores less than 42 corresponding to evening chronotypes and scores higher than 58 indicating morning chronotypes.

## **4. Pittsburgh Sleep Quality Index (PSQI)**

The evaluation of the participants’ sleep pattern quality was assessed with the PSQI scores (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989). The PSQI has 18 questions that yield seven component scores (sleep quality, sleep latency, duration, sleep efficiency, sleep disturbances, sleep medication use, and daytime dysfunction) rated from 0 (better) to 3 (worse). The total score, ranging from 0 (better) to 21 (worse), is the summation of all the component scores. Individuals with a PSQI total score  $\leq 5$  are characterized as good sleepers, whereas scores  $> 5$  are associated with poor sleep quality. According to its developers, the PSQI has a sensitivity of 89.6% and a specificity of 86.5% ( $\kappa = 0.75$ ,  $p < 0.001$ ), and an internal consistency  $\alpha = 0.83$  (Buysse et al., 1989).

## **5. Psychomotor Vigilance Test (PVT)**

Performance data were collected using the 3-minute PVT (Dinges & Powell, 1985) available on the AMI Motionloggers. PVT performance is not only affected by sleep loss, but it is also sensitive to circadian rhythmicity (Dinges et al., 1997; Doran, Van Dongen, & Dinges, 2001; Durmer & Dinges, 2005; Jewett, Dijk, Kronauer, & Dinges, 1999; Wyatt et al., 1997). The PVT is a simple reaction time test, where participants are required to press a response button as soon as the stimulus appears on the screen. The PVT has minor learning effects that can be reached in one to three trials (Dinges et al., 1997; Jewett et al., 1999; Kribbs & Dinges, 1994; Rosekind et al., 1994). The PVT interstimulus interval (ISI), defined as the period between the last response and the appearance of the next stimulus, ranges randomly from 2 to 10 seconds. The standard version of the PVT has a duration of 10 minutes (Loh, Lamond, Dorrian, Roach, & Dawson, 2004); however, shortened versions have also been used effectively to demonstrate sleep deprivation effects (Basner & Dinges, 2011; Loh et al., 2004). Since operational demands prevented the use of the 10-minute version, we used the 3-minute version of the PVT, which is included on the AMI Motionlogger actigraphs. The ISI ranged from 2 to 10 seconds, a red backlight appeared for one second and the word “PUSH” was the visual stimuli; the response time in milliseconds was then displayed to the participant.

## **6. Activity Logs**

All participants were asked to complete a daily activity log, documenting their daily routine in accordance with NSW categories. The logs covered 24-hour periods in 30-minute intervals.

## **D. PROCEDURES**

The study was approved by the Naval Postgraduate School Institutional Review Board. Personnel were briefed on the research protocol and study procedures. Those wishing to volunteer signed informed consent forms at the beginning of the study and received further training prior to being issued equipment for the study. Participants filled out the prestudy surveys upon receipt of their sleep watches and activity logs. All

participants were instructed to fill out their activity logs daily and, at a minimum, complete a PVT prior to and after their watchstanding period. Upon completion of the study, the participants returned their equipment and filled out an end-of-study survey.

## **E. VARIABLES**

Two independent variables, watch schedule and watch section, were used to compare the 3-hour on/9-hour off and 6-hour on/6-hour off watch schedules in terms of crew rest, sleep, and psychomotor vigilance performance. Dependent variables were daily rest and sleep amount (with and without naps within watch periods), number of sleep episodes per day, ESS scores, and PVT performance (as measured using mean reaction time [RT], mean response speed [1/RT], fastest 10% RT [i.e., 10<sup>th</sup> percentile of RT], slowest 10% of 1/RT [i.e., 10<sup>th</sup> percentile of 1/RT], percentage of 750ms, 600ms, 500ms and 355ms lapses, percentage of lapses and false starts, and percentage of false starts [FS]).

To assess work demands and rest opportunities, we compared the weekly amount of reported time with the time criteria of the NSW model activities (sleep, messing, personal time, free time, watch, work, training, service diversion, nonavailable time, productive work, and time on duty).

## **F. ANALYSIS**

### **1. Actigraphic Data Cleaning and Reduction Procedures**

The preparation of the actigraphic data for analysis included three steps. First, we determined the number of days of data available for each participant to examine the occurrence of gaps. Seventeen participants (13.8%) had fewer than five days of data and were omitted from the analysis. Then, we assessed the activity logs completed by each participant and verified the actigraphic data by using these logs. However, the main source of information for the sleep analysis was the actigraphic data, although sleep logs assisted in the determination of start and finish time of the sleep intervals. Based on this comparison, we manually identified the start and end times of sleep episodes in the actigraphic data. Imputation was applied only when: (a) there was a gap in actigraphic data within which the sleep log showed a sleep interval, or (b) the pattern of actigraphic

data, verified by the activity logs, was such to assure a confidence in the interpolation of a sleep interval. The criteria when considering imputation were the utility of the actigraphic data, the consistency in the pattern of activities over consecutive days, the amount of missing data, whether the participant was a watchstander, and the accuracy of the sleep log. Two crewmembers on the 3/9 schedule shifted their watch schedule for two or three days in the middle of the data collection period. We did not use their data from those days or from the day following their shift in schedule in the analysis. A large amount of data was missing on the first day of the study (3 December 2012); thus, our analysis used 4 December 2012 as the actual first day of the study.

Due to an operational commitment occurring on/about 15 December 2012, the ship was required to travel at high speeds, for a prolonged time, through rough seas. During this period, the motion of the ship was clearly evident in the actigraphic data, thereby contaminating the data for that time period. Therefore, all intervals after 15 December 2012 were omitted from the analysis. No observable interference due to ship motion was identified during the other days of the data collection period, although it should be noted that there is currently no objective method for isolating ship motion in actigraphic output. Using the methods described, an initial Excel spreadsheet of sleep intervals was developed that included data between 4 and 14 December 2012.

We focused on the rest/in-bed intervals (identified as DOWN intervals in the AMI software, and REST intervals in the Respironics software). The software provided the “Rest” time for each interval (also called time in-bed [TIB] in the literature); however, we did not use this naming convention, since some of the rest was obtained during watch. The software also provided the actigraphically evaluated Sleep amount within each Rest interval (also called Total Sleep Time [TST] in the literature). It should be noted that none of the participants reported a nap during watch periods, which was expected. For this reason, any naps within a watch period were based strictly on the pattern of activity that each participant demonstrated.

Of the 1,864 rest intervals, 80 (4.3%) were imputed, while 82 (4.4%) were naps identified during watch periods. The amount of rest and sleep for each day was calculated from 00:00 to 23:59.

## **2. PVT Data Cleaning and Reduction Procedures**

Psychomotor vigilance performance data were collected using the version of the PVT included on the AMI Motionloggers. The duration of each PVT trial was 3 minutes, with a minimum interstimulus interval (ISIMin) of 2 seconds and maximum interstimulus interval (ISIMax) of 10 seconds. As with the sleep analysis, tests taken on 3 December and after 15 December were omitted from the analysis. No imputation was applied in the PVT data.

PVT data were analyzed based on the metrics described by Basner and Dinges (2011) for individuals with chronic sleep deprivation. Specifically, a PVT response was scored as valid if RT was  $\geq 100$  milliseconds (ms). Responses without a stimulus or with RTs  $< 100$  ms were identified as false starts. Four categories of lapses were defined as RTs: equal to or greater than 355 ms, 500 ms, 600 ms, and 750 ms.

This data set, however, included some extraordinarily long reaction times of 10 seconds or more. Given that the PVT was not performed in controlled conditions, we postulate that these responses can be attributed predominantly to environmental factors acting as distractors (e.g., environmental noise, crewmembers in the same area, resumption of duties, etc.). For this reason, we omitted those responses with RTs  $\geq 10$  seconds from RT calculations, although we still denoted them as lapses and included them in the calculation of lapses ( $n = 78$  responses). With the ISI settings used, approximately 18 to 24 responses were expected in the 3-minute PVT. Therefore, trials with less than 10 responses were omitted from the analysis.

All PVT responses were aggregated by trial and then by participant. PVT performance metrics were analyzed between participants. No imputation was applied in the PVT data. PVT analysis was based on 959 trials.

## **3. Sleep Log Data Cleaning and Reduction Procedures for Workload Analysis**

Data from all sleep logs were screened for completeness and accuracy and were input into an Excel spreadsheet. Specifically, we assessed missing activity information or information that did not comply with the instructions for completing the sleep logs (e.g., adding activity codes not included in the instructions provided). Due to excessive

missing data, logs from 3 and 17 December were omitted from the analysis, which included data between 4 and 16 December 2012.

In some cases, participants included two activity codes in the same 30-minute interval ( $n = 102$ ; 0.18% of the 55,824 intervals). We interpreted these inputs as meaning that the participant spent time in both activities within the same 30-minute period. Without altering the amount of activity time within that day, we deleted the second activity from one cell and added it, as appropriate, to another cell with the same activity code. Two days of data were omitted from participant 3018 because of illness.

When deemed appropriate, data for days with missing activity were interpolated. The criteria for interpolation were (a) the accuracy of the sleep log, (b) the pattern of activities over consecutive days, (c) the length of missing data, (d) whether the participant was a watchstander, and (e) the existence of actigraphy data. Some logs were classified as inappropriate for interpolation because they did not correlate well with the actigraphic data. The pattern of activities was a critical criterion; that is, if the participants did not have a consistent daily pattern of activities, it was difficult to infer activities for missing days. Lastly, watchstanders had daily schedules that were more consistent and predictable. This consistency in activity patterns was not observed in nonwatchstanders; hence, we did not interpolate missing days of nonwatchstanders. Overall, we attempted to interpolate as needed, given the utility and accuracy of the available information sources, but we kept the interpolation rate as low as possible.

Entire day interpolation was applied to 59 days (5% of the 1,163), whereas partial interpolation was applied to five days that were missing, on average, 2.5 hours each. Overall, interpolation was applied to 2,344 30-minute intervals (4.2%).

#### **4. Analysis Roadmap**

Statistical analysis was conducted with the JMP Pro 9 statistical software package (SAS Institute; Cary, North Carolina). Descriptive statistical analysis was used to describe demographic characteristics of the population. Correlation analysis was conducted using nonparametric Spearman's rho. Levene's test was used for testing for equality of variances. Nonparametric methods were used for comparisons. These

included the Wilcoxon Rank Sum test and, for multiple comparisons, the Dunn method (for joint ranks accounting for group error rate).

Data are presented as mean (M)  $\pm$  standard deviation (SD) or median (MD) as appropriate. Significance level was set at  $p < 0.05$ . The nonparametric Wilcoxon Rank Sum test was used for comparisons. Correlation analysis was performed using the nonparametric Spearman's rho.

We initially performed a descriptive analysis of the entire data set, focusing on sleep intervals and daily rest/sleep amounts. Next, analysis was focused on the comparison between the 3/9 and 6/6 schedules.

### III. RESULTS

#### A. DEMOGRAPHICS

From the 122 crewmembers that initially volunteered, 52 were included in the watch schedule analysis, 41 (43.2%) in the 3/9 and 11 (11.6%) in the 6/6 watch schedules. Participants in the 3/9 stood watch between 00:01-03:00/12:00-15:00 (n = 6), 03:00-06:00/15:00-18:00 (n = 9), 06:00-09:00/18:00-21:00 (n = 8), and 09:00-12:00/21:00-23:59 (n = 18). Participants in the 6/6 stood watch between 00:01-06:00/12:00-18:00 (n = 7) and 06:00-12:00/18:00-23:59 (n = 4). All 6/6 participants were in the Operations (OPS) department, while the 3/9 participants were spread across the Weapons (WEPS), OPS, Engineering (ENG), and Supply departments. Appendix A provides a detailed description of participants by watch schedule, watch section, and actiwatch type. Table 1 shows the demographic information.

Table 1. Demographics.

Variable	Entire Data Set N = 122	Used for Analysis N = 52
Age, years, M ± SD	28 ± 5.96	29.1 ± 6.12
Gender, # males (%)	98 (80.3%)	40 (76.9%)
Pay grade, #		
Officers	28 (WO2-3, O1-5)	14 (WO2-3, O1-3)
Enlisted	94 (E2-8)	73 (E3-8)
Department, # All (Off.)		
CO – XO (not in a department)	2	–
Air Department	15 (6 officers)	–
Combat Systems	21 (3)	6 (4)
Engineering	32 (6)	19 (5)
Executive/Administration	3 (0)	–
Navigation	1 (0)	–
Operations	22 (5)	12 (4)
Supply	5 (1)	–
Weapons	20 (4)	1 (1)
Service, years, M ± SD	6.94 ± 5.55	7.15 ± 5.61
ME score, M ± SD	49.3 ± 8.05	47.3 ± 8.15
ME types, # (%)		
Definitely morning type	13 (10.7%)	–
Moderately morning type	1 (0.83%)	6 (11.5%)
Neither type	88 (72.70%)	34 (65.4%)
Moderately evening type	18 (14.90%)	12 (23.1%)
Definitely evening type	1 (0.83%)	–



## B. DESCRIPTIVE RESULTS FOR THE ENTIRE DATA SET

The average PSQI Global score for the entire data set was  $8.51 \pm 3.29$  (MD=8) ranging from 1 to 18. PSQI scores indicate that 91% of the participants were “poor sleepers” (PSQI score  $\geq 5$ ).

Actigraphic data were available from 95 participants over 1,864 sleep episodes. On average, each participant provided  $20.2 \pm 6.68$  sleep episodes (MD = 20), ranging from 9 to 44 sleep episodes per person. Analysis showed that participants slept, on average,  $1.77 \pm 0.598$  (MD = 1.75) episodes per day, ranging from 0.909 to 3.67 episodes. The average daily rest and sleep amount by participant is shown in Table 2.

Table 2. Daily rest and sleep amount by participant.

Sleep Intervals		Duration (hours)				
		Mean	Standard Deviation	Median	Minimum	Maximum
All	Rest	7.27	0.883	7.24	5.42	9.65
	Sleep	6.58	0.856	6.62	4.81	8.78
Only off-watch	Rest	7.20	0.903	7.17	4.65	9.65
	Sleep	6.53	0.873	6.60	4.07	8.78

Figures 2 and 3 show the frequency plots of daily rest and sleep amount in hours (all intervals included).




Figure 2. Daily rest amount in hours. Numbers on vertical bars represent the corresponding number of participants.




Figure 3. Daily sleep amount in hours. Numbers on vertical bars represent the corresponding number of participants.

It is notable that 95% of the participants received, on average, less than eight hours of sleep daily; 27% of the participants slept less than six hours per night. On average, participants experienced approximately 1.5 hours of sleep debt daily. Over the 11-day period of the research study, crewmembers accumulated an average of approximately 16 hours of sleep deficit. The average ESS score was  $10.6 \pm 3.70$  (MD = 10) ranging from 2 to 22. ESS scores indicate that 52 participants (42.6%) have excessive daytime sleepiness (i.e., ESS score > 10) (Johns, 1991).

### C. PRIMARY ANALYSIS

First, we assessed daily rest and sleep duration and compared them between schedules. From the actigraphic data, we determined that 10 (24.4%) participants on the 3/9 and 5 (45.5%) on the 6/6 schedules napped during watch. Analysis showed that the participants on the 6/6 schedule received less rest and sleep daily than participants on the 3/9 schedule, either when including all rest/sleep episodes or only those periods when sleep occurred during off-watch periods ( $p < 0.05$ ). When looking only at the off-watch sleep intervals, crewmembers on the 3/9 slept, on average,  $6.43 \pm 0.77$  hours, compared to  $5.66 \pm 0.79$  hours for those on the 6/6 schedule. When all sleep intervals included, crewmembers on the 3/9 slept, on average,  $6.46 \pm 0.77$  hours, compared to  $5.89 \pm 0.87$  hours for those on the 6/6 schedule. In short, the difference of approximately 46 minutes in daily sleep during off-watch periods was reduced to 34 minutes when including all rest/sleep intervals. Although napping during watch was identified in the actigraphic recordings of both watch schedules, napping on watch was more common on the 6/6

schedule. Consequently, crewmembers on the 6/6 partially compensated for their sleep debt by napping more during watch periods.

Including all sleep intervals, we assessed whether participants on the 6/6 schedule had more sleep episodes per day compared to those on the 3/9. Table 3 shows these results and indicates that there were significant differences between the 3/9 and 6/6 in mean values for all metrics. Figure 4 depicts daily sleep and sleep episodes per day.

Table 3. Daily rest/sleep amount in hours and number of sleep episodes per day by watch schedule.

Daily amount (hours)	3/9 M ± SD	6/6 M ± SD	Mean values Δ%	Sig. p val. <sup>A</sup>	Standard Deviation Δ%	Sig. p val. <sup>B</sup>
Daily Rest – all intervals	7.20 ± 0.71	6.49 ± 0.90	10.9%	0.007	-17.90%	0.622
Daily Rest – off-watch	7.17 ± 0.72	6.21 ± 0.78	16.5%	0.001	-4.88%	0.900
Daily Sleep – all intervals	6.46 ± 0.77	5.89 ± 0.87	11.2%	0.043	-11.10%	0.754
Daily Sleep – off-watch	6.43 ± 0.78	5.66 ± 0.79	15.8%	0.011	-2.44%	0.799
Sleep episodes per day	1.95 ± 0.50	2.39 ± 0.64	-18.4%	0.026	-21.90%	0.160

<sup>A</sup> Wilcoxon Rank Sum Test. <sup>B</sup> Levene’s test for equality of variances.




Figure 4. Daily sleep and number of sleep episodes per day by watch schedule.

Compared to participants on the 3/9 ( $10.7 \pm 3.70$ ), those watchstanders on the 6/6 had worse (i.e., larger) ESS scores ( $13.1 \pm 3.45$ ; Wilcoxon Rank Sum test:  $X^2(1) = 4.36$ ,  $p = 0.038$ ). Figure 5 depicts ESS scores by watch schedule.




Figure 5. Epworth Sleepiness Scale scores by watch schedule.

### 1. Comparison Between Watch Sections

We assessed changes in daily rest, sleep, and number of sleep episodes between various watch sections on the two watchstanding schedules. We identified a difference in daily rest (Dunn method,  $Z = 2.57$ ,  $p = 0.061$ ) and sleep (Dunn method,  $Z = 2.41$ ,  $p = 0.096$ ) between the 03:00-06:00 and the 06:00-09:00 sections on the 3/9 schedule. Figure 6 depicts daily sleep and number of sleep episodes per day, by the 3/9 sections. Vertical lines represent one standard deviation of daily sleep, including all sleep intervals or only those sleep intervals that occurred while participants were off watch.




Figure 6. Daily sleep and the number of sleep episodes per day in the 3/9 watch sections.

This diagram shows that the participants on the 3/9 schedule, with shifts occurring from midnight to 06:00 and from noon to 18:00, receive significantly more rest and sleep than their peers on the 06:00 to noon and 18:00 to midnight shifts. The same pattern is evident in the 6/6, but because of the small number of participants on the two shifts of the 6/6, statistical results are not reported. These results are shown in Table 4. Comparisons are based on Wilcoxon Rank Sum test.

Table 4. Daily rest/sleep amount in hours by watch shift.

Watch Schedule	Daily amount (hours)	1 <sup>A</sup> M ± SD	2 <sup>B</sup> M ± SD	X2(1)	Sig. p val. <sup>A</sup>
3/9	Rest – all intervals	7.48 ± 0.60	7.04 ± 0.73	3.90	0.048
	Rest – off watch	7.43 ± 0.64	7.02 ± 0.73	3.10	0.079
	Sleep – all intervals	6.82 ± 0.70	6.24 ± 0.74	4.69	0.030
	Sleep – off watch	6.78 ± 0.73	6.23 ± 0.74	3.80	0.051
6/6	Rest – all intervals	6.80 ± 1.00	5.95 ± 0.30	–	–
	Rest – off watch	6.37 ± 0.94	5.91 ± 0.27	–	–
	Sleep – all intervals	6.16 ± 1.00	5.43 ± 0.31	–	–
	Sleep – off watch	5.82 ± 0.96	5.41 ± 0.29	–	–

<sup>A</sup> For the 3/9, “1” refers to participants on the 00:00-03:00/12:00-15:00 and 03:00-06:00/15:00-18:00 shifts combined. For the 6/6, “1” refers to participants on the 00:00-06:00/12:00-18:00 shift. <sup>B</sup> For the 3/9, “2” refers to participants on the 06:00-09:00/18:00-21:00 and 09:00-12:00/21:00-24:00 shifts combined. For the 6/6, “2” refers to participants on the 06:00-12:00/18:00-24:00 shift.

Figure 7 shows daily rest and sleep, by watch schedule and section. Napping during watches is clearly evident for participants on the 6/6 watch schedule, with most naps occurring during the 00:00-06:00 watch period.




Figure 7. Daily rest and sleep, by watch schedule and shift.

## 2. Psychomotor Vigilance Performance

We also assessed the effect of watch schedule (3/9 versus 6/6) on PVT performance. Congruent with the rest/sleep results already described, the average values of the PVT were better for participants on the 3/9 compared to those on the 6/6, but not at statistically significant levels. However, the two watch schedules differed significantly in variability; participants on the 6/6 had larger variability than those on the 3/9 for 11 of the 13 PVT metrics used ( $p < 0.05$ ). These results are shown in Table 5.

Table 5. Comparison of PVT metrics between the 3/9 and 6/6 watch schedules.

Variable	3/9 M ± SD	6/6 M ± SD	Mean values Δ%	Sig. p value <sup>A</sup>	Standard Deviation Δ%	Sig. p value <sup>B</sup>
Mean RT, [ms]	323 ± 66.9	372 ± 135	-13.20%	0.407	-50.4%	0.009
Mean 1/RT	3.95 ± 0.524	3.67 ± 0.928	7.63%	0.293	-43.5%	0.016
Fastest 10% RT, [ms]	196 ± 28.0	217 ± 52.7	-9.68%	0.275	-46.9%	0.019
Slowest 10% 1/RT	2.43 ± 0.469	2.18 ± 0.743	11.50%	0.374	-36.9%	0.069
False Starts (FS), %	2.0 ± 1.59	2.23 ± 2.10	-10.30%	0.936	-24.3%	0.474
Lapses 750ms, %	3.74 ± 2.60	5.70 ± 5.38	-35.40%	0.492	-51.7%	0.007
Lapses 600ms, %	5.30 ± 3.06	8.39 ± 7.19	-36.8%	0.518	-57.4%	< 0.001
Lapses 500ms, %	7.54 ± 4.40	11.9 ± 9.57	-36.60%	0.332	-54.0%	< 0.001
Lapses 355ms, %	17.0 ± 9.74	26.8 ± 18.5	-36.60%	0.210	-47.4%	0.005
Lapses 750ms+FS, %	5.74 ± 3.74	7.98 ± 4.86	-28.10%	0.210	-23.1%	0.591
Lapses 600ms+FS, %	7.30 ± 4.03	10.7 ± 6.50	-31.80%	0.258	-38.0%	0.068
Lapses 500ms+FS, %	9.54 ± 5.09	14.2 ± 8.64	-32.80%	0.293	-41.1%	0.014
Lapses 355ms+FS, %	19.0 ± 9.78	29.1 ± 17.2	-34.70%	0.182	-43.1%	0.008

<sup>A</sup> Wilcoxon Rank Sum Test. <sup>B</sup> Levene's test for equality of variances.



Figure 8 shows the aforementioned differences in variability between watch schedules.




Figure 8. Differences in standard deviations in PVT metrics between the 3/9 and 6/6 watch schedules.

### 3. Posttest Questionnaires

At the completion of the study, participants (N = 122) rated seven watch schedules (5/10, 5/15, 6/6, 3/9, 12/12, 6/12, and 6/18). They were asked to respond to the question “Compared to my current schedule, the [watch schedule] is ...” using a 6-level Likert scale (Worse “1,” Same as “2,” Better “3,” Never stood the [watch schedule] watch before “4,” Standing [watch schedule] watch now “5,” No answer “6”). Results show that the participants evaluate the 6/6 as being the worst schedule and the 3/9 and 6/18 as the two best. Figure 9 shows the integrated results by watch schedule.




Figure 9. Subjective evaluations of watch schedules.

#### 4. Work Patterns

Next, we assessed work patterns in terms of NSWW compliance and the association between work patterns and watch schedule. Crewmembers on the 6/6 watch schedule work 30% more than the NSWW criterion (105 hours compared to 81 hours) and stand watch 60% more than the criterion (91 hours compared to 56 hours). We found that participants on the 6/6 watch schedule experience a 15-hour workday. Furthermore, crewmembers on the 6/6 schedule indicate spending 460% more time on Service Diversion than the NSWW criterion and over twice as much as participants on the 3/9 (+225%). These results are shown in Table 6.

Table 6. Work patterns by watch schedule.

Activity (criterion)	3/9 (M ± SD)	6/6 (M ± SD)	Watch schedule comparison p-value
Non-available time (87 hours)	82.6 ± 13.6**	63.0 ± 4.84**	< 0.001
Sleep (56 hours)	46.6 ± 7.10***	42.2 ± 8.30**	0.167
Messing (14 hours)	10.4 ± 4.00***	8.24 ± 2.70**	0.157
Personal time (14 hours)	24.4 ± 13.0***	12.5 ± 5.77	< 0.001
Free time (3 hours)	1.26 ± 2.80***	0	
On Duty (81 hours)	85.4 ± 13.6**	105 ± 4.84**	< 0.001
Productive Work (70 hours)	58.2 ± 14.4***	91.4 ± 3.84**	< 0.001
Watch (56 hours)	48.7 ± 8.60***	90.0 ± 2.20**	< 0.001
Work (14 hours)	9.55 ± 12.2**	1.32 ± 4.00**	0.028
Training (7 hours)	4.84 ± 5.10**	0.695 ± 0.900**	0.003
Service diversion (4 hours)	22.4 ± 15.9***	13.0 ± 6.70**	0.102

Note: Different than the criterion: \* p < 0.05; \*\* p < 0.01; \*\*\* p < 0.001

Service Diversion activities are responsible for dramatically reducing sleep opportunities for participants on the 6/6 watch schedule. This result was further assessed by a stepwise regression analysis. Initially, we included all activity groups as probable factors for consideration. The stepwise analysis showed that only the Service Diversion group was associated with sleep time ( $p = 0.05$ ). Based on this result, the corresponding regression model showed that Service Diversion alone explained 62% of the observed variability in the sleep time of participants on the 6/6,  $R^2_{\text{adj}}=0.575$ ,  $F(1, 8) = 13.2$ ,  $p = 0.007$ . Stepwise analysis of the sleep patterns for participants on the 3/9 did not exclude any of the seven groups of activity. The corresponding multiple regression model explained 86% of the observed variability of sleep time of participants on the 3/9,  $R^2_{\text{adj}} = 0.832$ ,  $F(7, 36) = 31.4$ ,  $p < 0.001$ .

The considerable differences in distributions of duty time, productive work, watch, and sleep between participants on the 3/9 and 6/6 watch schedules are clearly depicted in Figures 10-16. Vertical axes show the percentage of participants on each watch section. Each time bin represents a 30-minute period.




Figure 10. Work and sleep patterns in the 00:00 – 03:00 and 12:00 – 15:00 section of the 3/9 watch schedule.




Figure 11. Work and sleep patterns in the 03:00 – 06:00 and 15:00 – 18:00 section of the 3/9 watch schedule.




Figure 12. Work and sleep patterns in the 06:00 – 09:00 and 18:00 – 21:00 section of the 3/9 watch schedule.




Figure 13. Work and sleep patterns in the 09:00 – 12:00 and 21:00 – 23:59 section of the 3/9 watch schedule.




Figure 14. Work and sleep patterns in the 00:00 – 06:00 and 12:00 – 18:00 section of the 6/6 watch schedule.




Figure 15. Work and sleep patterns in the 06:00 – 12:00 and 18:00 – 23:59 section of the 6/6 watch schedule.





Figure 16. Polar diagrams of typical 24-hour days in the 3/9 (left diagram) and 6/6 watch schedules (right diagram).

## IV. DISCUSSION

Our results show that from a human-centered perspective, the 3/9 watch schedule is better than the 6/6 schedule in terms of patterns of work, rest and sleep, subjective levels of fatigue, psychomotor vigilance performance, and acceptance by the participants. Although crewmembers were sleep deprived on both watch schedules, crewmembers on the 3/9 received more sleep than their peers in the 6/6,  $6.46 \pm 0.77$  hours and  $5.89 \pm 0.87$  hours, respectively. The 3/9 schedule was also better than the 6/6 in terms of the distribution of sleep episodes. Specifically, crewmembers on the 3/9 received more night sleep, whereas the crewmembers on the 6/6 had to sleep during the day to compensate for their sleep loss at night.

Actigraphic analysis also showed that 24.4% participants on the 3/9 schedule and 45.5% on the 6/6 schedule napped occasionally during their watch. Napping was more pronounced in crewmembers on the 6/6 schedule during the night watches, which fits with earlier research showing that involuntary sleep is more commonly experienced on night shifts, with 7% to 20% of the personnel reporting falling asleep during night work (Åkerstedt et al., 2002; Åkerstedt & Wright, 2009). However, whether a specific watch location affords involuntary napping depends on the type of watch duties assigned. Therefore, while severe sleep debt and its concomitant need for napping almost certainly exist in other watch stations, it is not feasible for personnel to nap because of their assigned duties. The issue of falling asleep while on watch is critical because it indicates how, in extreme sleep debt, the individual uses every possible opportunity to compensate for the accumulated sleep debt. Future efforts should investigate whether napping during watch is a viable operational measure to ameliorate sleep deprivation, in conjunction with the specific duties of each watch location. We should note that the identification of naps was based solely on actigraphic activity patterns. None of the activity logs reported a nap within a watch period, probably due to the controversial nature of such a statement from a military member. Yet, our approach for identifying periods of low activity as naps was conservative; we identified naps only when the activity change was clear and distinct.

These results clearly show that the 3/9 schedule is better than the 6/6 from a sleep hygiene perspective. Not surprisingly, the impact of watch schedule on sleep hygiene is



evident in the ESS scores with participants on the 3/9 showing decreased daytime sleepiness compared to their peers on the 6/6.

In terms of work demands, crewmembers on the 6/6 have considerably long workdays (on average, 15 hours on duty), which corresponds to approximately 30% more time on duty than the NSWW criterion (105 hours compared to 81 on a weekly basis). These results are generally congruent with earlier research on U.S. Navy ships in which over 50% of the participants worked more than 95 hours per week, approximately 13.6 hours per day (Haynes, 2007). This amount of daily work far exceeds the recommended work hours, especially when considering that work hours in excess of 8 to 12 hours daily could threaten crew members' ability to perform safe operations (Comperatore, Kingsley, Kirby, & Rivera, 2001). Our findings should also be considered in light of the work hour regulations specified in Title 46 of the United States Code, 2006 Edition, Supplement 5, Section 8104, which states that except in an emergency, licensed individuals cannot be required to work more than 12 of 24 hours at sea ("United States Code," 2006). It further states that on oil tankers, licensed individuals or seamen may not be permitted to work more than 15 hours in any 24-hour period, or more than 36 hours in any 72-hour period—except in an emergency or a drill. It should be noted that the term "work" in the U.S. Code corresponds to the NSWW "time on duty," which includes any administrative duties.

Given the differences in sleep hygiene and work demands between the two watch schedules, the psychomotor vigilance performance results are not surprising. The two schedules differed significantly in variability. Compared to their 3/9 counterparts, personnel on the 6/6 schedule showed greater variability in 11 of the 13 PVT metrics we used ( $p < 0.05$ ). Furthermore, the average values of the PVT metrics were better for personnel on the 3/9 compared to the 6/6, although not at statistically significant levels. The fact that the personnel on the 6/6 schedule had significantly increased variability in psychomotor performance is a major concern in the operational environment. Miller (2006) highlighted performance variability as a primary hallmark of human fatigue, even though performance variability, in itself, has received little attention as a measure of performance impairment (J. C. Miller, 2013a). He attributed this variability in

performance to large amplitude, moment-to-moment fluctuations in attention associated with fatigue.

The findings of this study are integrated in Table 7, which assesses the overall utility of the two watch schedules under focus (3/9, 6/6) based on their ranking. A score of 1 is considered the best rank (e.g., more sleep, more rest, fewer sleep episodes per day yielding less fragmented sleep, lower ESS score, etc.). Sleeping on watch is a negative phenomenon; thus, a ranking of 1 refers to the schedule with less napping, whereas the lower rank denotes the schedule with the largest percentage of individuals sleeping on watch. The overall ranking is based on the comparison between 3/9 and 6/6 schedules. The ranking assigned for each factor shows a general pattern of differences between schedules that support any inferential statistics already presented.

Table 7. Comparison ranks between watch schedules.

Watch Schedule	3/9 n = 45	6/6 n = 11
Daily Rest amount – All intervals	1	2
Daily Sleep amount – All intervals	1	2
Affordability of sleep in off-watch periods	1	2
Napping in watches	1	2
Sleep episodes per day	Inconclusive	
ESS	1	2
PVT performance	1	2
Subjective assessment of the watch schedule	1	2
Work patterns	1	2
Overall	1	2

From a human-centered perspective, the pattern of differences between 3/9 and 6/6, combined with the overall ranking, show clearly that the 3/9 is a better watch schedule compared to the 6/6 in terms of rest and sleep, subjective levels of fatigue, psychomotor performance, and acceptance from the participants. This conclusion aligns with previous research (Paul et al., 2010), which showed that the FAST predicted effectiveness of the 6/6 is worse compared to the 8/16, the 4/8, and the 8on-4off-4on-8off watch schedules. The 6/6 schedule should be avoided when alternative circadian-aligned watch schedules can be used.

A critical constraint when addressing optimization of watch schedules, however, is the availability of qualified watchstanders. The two watch-standing schedules we compared have vastly different characteristics in terms of personnel needed to implement them. The 3/9 schedule is a 4-section watchbill in which individuals stand watch for only six hours per day; the 6/6 has only two sections with individuals standing watch for 12 hours each day; the 6/6 theoretically takes only half the crewmembers needed by the 3/9 schedule. This perspective, though, oversimplifies the problem of optimized shiftwork to a simple enumeration of the minimum number of people needed. Such approaches have been used in the past and inevitably lead to watch systems with increased sleep deprivation, fatigue, and circadian desynchrony (Miller, 2013a, 2013b).

#### **A. SOME THOUGHTS ON WORK DEMANDS AND REST AFFORDABILITY**

Our results show that while underway, personnel receive much less sleep than the eight hours per day that is widely accepted as normal. This result agrees with earlier research findings conducted on Navy ships (Green, 2009; Haynes, 2007; Mason, 2009; Miller, Matsangas, & Kenney, 2012); however, we have also found that personnel spend a considerable amount of time for their personal needs. The question raised is why don't crewmembers spend more of their personal time sleeping in order to ameliorate their accumulated sleep debt?

One plausible explanation lies in the characteristics of the naval environment and life at sea for extended periods of time. A crew onboard a ship works in a confined environment, with limited opportunities for privacy; their daily schedule continues around the clock and they continue to work without weekends and holidays, away from family and friends until the ship returns to port. Especially evident under these grueling work conditions, is that organizational structure plays a critical role in the psychological health of the crew members (e.g., by providing coworker support and the social interactions that are vital for good morale and mental health) (Maslow, 1943; Taormina, 1997). Although this study was not focused on this issue *per se*, future efforts should investigate how psychological health is affected by increased workload combined with the unique stresses presented by the shipboard environment.

Educating crewmembers on healthy sleep habits is also essential. If personnel were more attuned to the consequences of sleep deprivation on their performance and safety, it is likely that they would try to manage their schedules more wisely. Hence, sleep hygiene training could be beneficial as a fatigue countermeasure.

Another issue of interest is the considerable amount of time spent in Service Diversion activities. Reducing the demand for these activities may free up time for other needs. Some researchers have noted that reducing administrative tasks is a necessary and effective measure for reducing fatigue on naval vessels (Houtman et al., 2005). However, some of the Service Diversion activities may be based on organizational regulations, and thus, may not fall under the command authority of the commanding officer of the ship.

Lastly, the NSW model should be revised to include adequate time for rest, part of which is the actual time set aside for sleep. Rest time involves more than just the time dedicated for sleep since it takes some time to decompress, fall asleep and awaken. The International Maritime Organization (IMO) has proposed a minimum of 10 hours of **rest** during any 24-hour period and 77 hours of rest for any 7-day period (International Maritime Organization, 2010).

## **B. CAVEATS**

This study has a number of caveats. First, the participants were volunteers, performing their normal daily duties; there was no randomization in the assignment to watch schedule and hence the study is quasi-experimental in nature. Second, all the participants on the 6/6 watch schedule were from the Operations Department. Third, some groups in this analysis had fewer participants than originally planned, resulting in unequal cell sizes and making analysis challenging.

THIS PAGE INTENTIONALLY LEFT BLANK

## APPENDIX. PARTICIPANT INFORMATION

Table 8 shows participants by watch schedule, watch section, and actiwatch type.

Table 8. Participants by watch schedule, watch section, and actiwatch type.

Watch Schedule	Watch Section	AMI	PR
3/9 (n = 41)	00:00 - 03:00	n = 5: 3001, 3007, 3025, 3032, 3057	n = 1: 3122
	03:00 - 06:00	n = 5: 3036, 3040, 3054, 3061, 3069	n = 4: 3102, 3110, 3112, 3113
	06:00 - 09:00	n = 6: 3006, 3009, 3016, 3022, 3027, 3090	n = 2: 3101, 3109
	09:00 - 12:00	n = 11: 3002, 3017, 3021, 3043, 3055, 3063, 3074, 3078, 3079, 3080, 3093	n = 7: 3104, 3108, 3111, 3117, 3118, 3119, 3131
4/8 (n = 8)	00:00 - 04:00	n = 2: 3013, 3146	
	04:00 - 08:00	n = 1: 3082	
	08:00 - 12:00	n = 5: 3056, 3064, 3067, 3073, 3092	
6/6 (n = 11)	00:00 - 06:00	n = 6: 3011, 3018, 3026, 3028, 3030, 3084	n = 1: 3134
	06:00 - 12:00	n = 4: 3015, 3035, 3095, 3096	
6/12 (n = 3)	Shifting schedule, no fixed sections	n = 3: 3045, 3089, 3100	
6/18 (n = 4)	06:00 - 12:00	–	
	12:00 - 18:00	n = 2: 3083, 3085	
	18:00 - 23:59	n = 2: 3020, 3047	
12/12 (n = 5)	00:00 - 12:00	n = 2: 3031, 3097	
	12:00 - 23:59	n = 3: 3023, 3058, 3077	
Nonwatch standers (NWS) (n = 23)		n = 16: 3005, 3008, 3033, 3034, 3041, 3042, 3051, 3059, 3060, 3062, 3065, 3070, 3071, 3076, 3087, 3091	n = 7: 3120, 3124, 3125, 3127, 3128, 3129, 3153

THIS PAGE INTENTIONALLY LEFT BLANK

## LIST OF REFERENCES

- Åkerstedt, T. (2003). Shift work and disturbed sleep/wakefulness. *Occupational Medicine*, *53*, 89–94.
- Åkerstedt, T., Knutsson, A., Westerholm, P., Theorell, T., Alfredsson, L., & Kecklund, G. (2002). Work organisation and unintentional sleep: Results from the WOLF study. *Occupational and Environmental Medicine*, *59*, 595–600.
- Åkerstedt, T., & Wright, K. P. (2009). Sleep loss and fatigue in shift work and shift work disorder. *Sleep Medicine Clinics*, *4*, 257–271.
- Albertsen, K., Rafnsdóttir, G. L., Grimsmo, A., Tómasson, K., & Kauppinen, K. (2008). Workhours and worklife balance. *Scandinavian Journal of Work, Environment & Health, Supplements*, *5*, 14–21.
- Arendt, J., Middleton, B., Williams, P., Francis, G., & Luke, C. (2006). Sleep and circadian phase in a ship's crew. *Journal of Biological Rhythms*, *21*, 214–221.
- Arnedt, J. T., Owens, J., Crouch, M., Stahl, J., & Carskadon, M. A. (2005). Neurobehavioral performance of residents after heavy night call vs. after alcohol ingestion. *JAMA*, *294*, 1025–1033.
- Basner, M., & Dinges, D. F. (2011). Maximizing sensitivity of the Psychomotor Vigilance Test (PVT) to sleep loss. *Sleep*, *34*(5), 581–591.
- Belenky, G., Wesensten, N. J., Thorne, D. R., Thomas, M. L., Sing, H. C., Redmond, D. P., . . . Balkin, T. J. (2003). Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: A sleep dose-response study. *Journal of Sleep Research*, *12*, 1–12.
- Burch, J. B., Yost, M. G., Johnson, W., & Allen, E. (2005). Melatonin, sleep, and shift work adaptation. *Journal of Occupational and Environmental Medicine*, *47*(9), 893–901.
- Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Journal of Psychiatric Research*, *28*(2), 193–213.
- Colquhoun, W. P., Blake, M. J. F., & Edwards, R. S. (1969a). Experimental studies of shift work III: Stabilized 12-hour shift systems. *Ergonomics*, *12*, 865–882.
- Colquhoun, W. P., Blake, M. J. F., & Edwards, R. S. (1969b). Experimental studies of shift-work I: A comparison of “rotating” and “stabilized” 4-hour shift systems. *Ergonomics*, *11*, 437–453.



- Colquhoun, W. P., & Folkard, S. (1985). Scheduling watches at sea. In S. Folkard & T. H. Monk (Eds.), *Hours of work: Temporal factors in work-scheduling* (pp. 253–260). New York: John Wiley and Sons.
- Comperatore, C. A., Kingsley, L., Kirby, A., & Rivera, P. K. (2001). Management of endurance risk factors: A guide for deep draft vessels. Groton, CT: U.S. Coast Guard Research and Development Center.
- Cruz, C., Boquet, A., Detwiller, C., & Nesthus, T. (2003). Clockwise and counterclockwise rotating shifts: effects on vigilance and performance. *Aviation Space and Environmental Medicine*, 74(6), 606–614.
- Department of the Navy. (2007). Navy total force manpower policies procedures. Washington, D.C.: Department of the Navy.
- Department of the Navy. (2012). Standard organization and regulations of the U.S. Navy - OPNAV Instruction 3120.32D. Washington, D.C.: Department of the Navy.
- Dijk, D. J., & Czeisler, C. A. (1995). Contribution of the circadian pacemaker and the sleep homeostat to sleep propensity, sleep structure, electroencephalographic slow waves, and sleep spindle activity in humans *The Journal of Neuroscience*, 15, 3526–3538.
- Dijk, D. J., & Edgar, D. M. (1999). Circadian and homeostatic control of wakefulness and sleep. In F. W. Turek & P. C. Zee (Eds.), *Regulation of sleep and wakefulness* (pp. 111–147). New York: Marcel Dekker.
- Dinges, D. F., Pack, F., Williams, K., Gillen, K. A., Powell, J. W., Ott, G. E., . . . Pack, A. I. (1997). Cumulative sleepiness, mood disturbance, and psychomotor vigilance performance decrements during a week of sleep restricted to 4-5 hours per night. *Sleep*, 20(4), 267–277.
- Dinges, D. F., & Powell, J. W. (1985). Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *Behavior Research Methods, Instruments, & Computers*, 17(6), 652–655.
- Doran, S. M., Van Dongen, H. P. A., & Dinges, D. F. (2001). Sustained attention performance during sleep deprivation: Evidence of state instability. *Archives Italiennes de Biologie*, 139(3), 253–267.
- Drake, C. L., & Wright, K. P. (2011). Shift work, shift-work disorder, and jet lag. In M. H. Kryger, T. Roth, & W. C. Dement (Eds.), *Principles and Practices of Sleep Medicine* (5th ed., pp. 784–798). St. Louis, MO: Elsevier Saunders.
- Durmer, J. S., & Dinges, D. F. (2005). Neurocognitive consequences of sleep deprivation. *Seminars in Neurology*, 25(1), 117–129.

- Folkard, S. (1992). Is there a “best compromise” shift system? *Ergonomics*, 35, 1453–1463.
- Folkard, S., Lombardi, D. A., & Spencer, M. B. (2006). Estimating the circadian rhythm in the risk of occupational injuries and accidents. *Chronobiology International*, 23(6), 1181–1192.
- Folkard, S., & Tucker, P. (2003). Shift work, safety and productivity. *Occupational Medicine*, 53, 95–101.
- Goh, V. H., Tong, T. Y., Lim, C. L., Low, E. C., & Lee, L. K. (2000). Circadian disturbances after night-shift work onboard a naval ship. *Military Medicine*, 165(2), 101–105.
- Green, K. Y. (2009). *A comparative analysis between the Navy standard workweek and the actual work/rest patterns of sailors aboard U.S. Navy frigates* (Master’s thesis). Monterey, CA: Naval Postgraduate School.
- Hakola, T., & Härmä, M. (2001). Evaluation of a fast forward rotating shift schedule in the steel industry with a special focus on ageing and sleep. *Journal of Human Ergology*, 30(1–2), 315–319.
- Harrington, J. M. (2001). Health effects of shift work and extended hours of work. *Occupational and Environmental Medicine*, 58, 68–72.
- Haynes, L. E. (2007). *A comparison between the Navy standard workweek and the actual work and rest patterns of U.S. Navy sailors* (Master’s thesis). Monterey, CA: Naval Postgraduate School.
- Hockey, R. (Ed.). (1983). *Stress and Fatigue in Human Performance*: John Wiley & Sons.
- Horne, J. A., & Östberg, O. (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International Journal of Chronobiology*, 4, 97–110.
- Houtman, I., Miedema, M., Jettinghoff, K., Starren, A., Heinrich, J., Gort, J., . . . Wubbolts, S. (2005). *Fatigue in the shipping industry*. Hoofddorp, Netherlands: TNO Work and Employment.
- Howarth, H. D., Pratt, J., & Tepas, D. I. (1999). Do maritime crew members have sleep disturbances? *International Journal of Occupational and Environmental Health*, 5(2), 95–100.
- Hursh, S. R., Redmond, D. P., Johnson, M. L., Thorne, D. R., Belenky, G., Balkin, T. J., . . . Eddy, D. R. (2004). Fatigue models for applied research in warfighting. *Aviation Space and Environmental Medicine*, 75(3 Suppl), A44–A53; discussion A54–A60.

- International Maritime Organization. (2010). Standards of training, certification and watchkeeping for seafarers: IMO.
- Jewett, M. E., Dijk, D. J., Kronauer, R. E., & Dinges, D. F. (1999). Dose-response relationship between sleep duration and human psychomotor vigilance and subjective alertness. *Sleep*, 22(2), 171–179.
- Johns, M. W. (1991). A new method for measuring daytime sleepiness: The Epworth Sleepiness Scale. *Sleep*, 14, 540–545.
- Johns, M. W. (1992). Reliability and factor analysis of the Epworth Sleepiness Scale. *Sleep*, 15(4), 376–381.
- Knutsson, A. (1989). Shift work and coronary heart disease. *Scandinavian Journal of Social Medicine*, 44(Supplementum), 1–36.
- Knutsson, A. (2003). Health disorders of shift workers. *Occupational Medicine*, 53, 103–108.
- Kribbs, N. B., & Dinges, D. F. (1994). Vigilance decrement and sleepiness. In J. R. Harsh & R. D. Ogilvie (Eds.), *Sleep Onset Mechanisms* (pp. 113–125). Washington, D.C.: American Psychological Association.
- Kronauer, R. E., & Stone, B. M. (2004). Commentary on fatigue models for applied research in warfighting. *Aviation Space and Environmental Medicine*, 75(3, Suppl.), A54–A56.
- Loh, S., Lamond, N., Dorrian, J., Roach, G., & Dawson, D. (2004). The validity of psychomotor vigilance tasks of less than 10-minute duration. *Behavior Research Methods, Instruments, & Computers*, 36(2), 339–346.
- Maslow, A. H. (1943). A theory of human motivation. *Psychological Review*, 50, 370–396.
- Mason, D. R. (2009). *A comparative analysis between the Navy standard workweek and the work/ rest patterns of Sailors aboard U.S. Navy cruisers* (Master's thesis). Monterey, CA: Naval Postgraduate School.
- Meltzer, L. J., Walsh, C. M., Traylor, J., & Westin, A. M. L. (2012). Direct comparison of two new actigraphs and polysomnography in children and adolescents. *Sleep*, 35, 159–166.
- Miller, J. C. (2006). Fundamentals of shiftwork scheduling. Brooks City-Base, TX: Air Force Research Laboratory.
- Miller, J. C. (2013a). *Fundamentals of Shiftwork Scheduling, 3rd Edition: Fixing Stupid*.
- Miller, J. C. (2013b). *Shiftwork: An annotated bibliography*.

- Miller, N. L., Matsangas, P., & Kenney, A. (2012). The role of sleep in the military: implications for training and operational effectiveness. In J. H. Laurence & M. D. Matthews (Eds.), *The Oxford Handbook of Military Psychology* (pp. 262–281): Oxford University Press.
- Monk, T. H. (1986). Advantages and disadvantages of rapidly rotating shift schedules - A circadian viewpoint. *Human Factors*, 28, 553–557.
- Monk, T. H. (1990). The relationship of chronobiology to sleep schedules and performance demands. *Work & Stress*, 4(3), 227–236.
- Monk, T. H., Buysse, D. J., Billy, B. D., Fletcher, M. E., Kennedy, K. S., Begley, A. E., . . . Beach, S. R. (2013). Shiftworkers report worse sleep than day workers, even in retirement. *Journal of Sleep Research*, 22, 201–208.
- Osborn, C. M. (2004). *An analysis of the effectiveness of a new watchstanding schedule for U.S. submariners* (Master's thesis). Monterey, CA: Naval Postgraduate School.
- Paul, M. A., Ebisuzaki, D., McHarg, J., Hursh, S. R., & Miller, J. C. (2012). An assessment of some watch schedule variants used in Canadian patrol frigates. Toronto, Canada: Defence Research and Development.
- Paul, M. A., Hursh, S. R., & Miller, J. C. (2010). Alternative submarine watch systems: Recommendation for a new CF submarine watch schedule: Defence Research and Development Canada (DRDC).
- Rosa, R. R., Bonnet, M. H., & Cole, L. L. (1998). Work schedule and task factors in upper-extremity fatigue. *Human Factors*, 40(1), 150–158.
- Rosekind, M. R., Graeber, R. C., Dinges, D. F., Connel, L. J., Rountree, M. S., & Spinweber, C. L. (1994). Crew factors in flight operations IX: Effects of planned cockpit rest on crew performance and alertness in long-haul operations. Moffett Field, CA: NASA Ames Research Center.
- Rutenfranz, J., Colquhoun, W. P., Knauth, P., & Ghata, J. N. (1977). Biomedical and psychosocial aspects of shift work. *Scandinavian Journal of Work, Environment & Health*, 3(4), 165–182.
- Rutenfranz, J., Plett, R., Knauth, P., Condon, R., De Vol, D., Fletcher, N., . . . Colquhoun, W. P. (1988). Work at sea: A study of sleep, and circadian rhythms in physiological and psychological functions, in watchkeepers on merchant vessels. *International Archives of Occupational and Environmental Health*, 60(5), 331–339.
- Sallinen, M., & Kecklund, G. (2010). Shift work, sleep, and sleepiness - Differences between shift schedules and systems. *Scandinavian Journal of Work, Environment & Health*, 36, 121–133.

- Stevens, R. G., Hansen, J., Costa, G., Haus, E., Kauppinen, T., Aronson, K. J., . . . Straif, K. (2011). Considerations of circadian impact for defining “shift work” in cancer studies: IARC Working Group Report. *Occupational and Environmental Medicine*, 68, 154–162.
- Stolgitis, W. C. (1969). *The effects of sleep loss and demanding work/rest cycles: An analysis of the traditional Navy watch system and a proposed alternative* (Master’s thesis). Monterey, CA: Naval Postgraduate School.
- Taormina, R. J. (1997). Organizational socialization: A multidomain, continuous process model. *International Journal of Selection and Assessment*, 5(1), 29–47.
- United States Code, Supplement 5, Title 46 § 8104 (2006).
- Viitasalo, K., Kuosma, E., Laitinen, J., & Härmä, M. (2008). Effects of shift rotation and the flexibility of a shift system on daytime alertness and cardiovascular risk factors. *Scandinavian Journal of Work, Environment & Health*, 34(3), 198–205.
- Wagstaff, A. S., & Sigstad Lie, J.-A. (2011). Shift and night work and long working hours – A systematic review of safety implications. *Scandinavian Journal of Work, Environment & Health*, 37(3), 173–185.
- Wang, X.-S., Armstrong, M. E. G., Cairns, B. J., Key, T. J., & Travis, R. C. (2011). Shift work and chronic disease: The epidemiological evidence. *Occupational Medicine*, 61, 78–89.
- Wedderburn, A. A. I. (1992). How fast should the night shift rotate? A rejoinder. *Ergonomics*, 35, 1447–1451.
- Wilkinson, R. T. (1992). How fast should the night shift rotate? *Ergonomics*, 35, 1425–1446.
- Wyatt, J. K., Dijk, D. J., Ronda, J. M., Jewett, M. E., Powell, J. W., Dinges, D. F., & Czeisler, C. A. (1997). Interaction of circadian- and sleep/wake homeostatic-processes modulate psychomotor vigilance test (PVT) performance (Abstract). *Sleep Research*, 26, 759.
- Yamada, C., Tachibana, M., & Kuriyama, K. (1988). Neurochemical changes in the cholinergic system of the rat lateral vestibular nucleus following hemilabyrinthectomy. *Arch Otorhinolaryngol*, 245(4), 197–202.
- Yokeley, M. T. (2012). *Effects of sleep deprivation on U.S. Navy surface ship watchstander performance using alternative watch schedules* (Master’s thesis). Monterey, CA: Naval Postgraduate School.

## INITIAL DISTRIBUTION LIST

1. Defense Technical Information Center  
Ft. Belvoir, Virginia
  
2. Dudley Knox Library  
Naval Postgraduate School  
Monterey, California
  
3. Research Sponsored Programs Office, Code 41  
Naval Postgraduate School  
Monterey, California
  
4. Richard Mastowski (Technical Editor).....1  
Graduate School of Operational and Information Sciences (GSOIS)  
Naval Postgraduate School  
Monterey, California
  
5. Nita Lewis Shattuck.....1  
Operations Research Department  
Graduate School of Operational and Information Sciences (GSOIS)  
Naval Postgraduate School  
Monterey, California
  
6. Panagiotis Matsangas.....1  
Operations Research Department  
Graduate School of Operational and Information Sciences (GSOIS)  
Naval Postgraduate School  
Monterey, California