


14th Symposium on Integrated Observing and Assimilation Systems for the
Atmosphere, Oceans, and Land Surface (IOAS-AOLS), AMS
Atlanta, January 17-21, 2010

Eddy and Chlorophyll-a Structure in the Kuroshio Extension Detected from Altimeter and SeaWiFS

Peter C. Chu and Yu-heng Kuo
Naval Postgraduate School
Monterey, CA 93943

pcchu@nps.edu; <http://faculty.nps.edu/pcchu>

Mean Surface Dynamic Height 170 cm Contour → Kuroshio Extension (KE) (Qiu & Chen 2005 JPO)



Theories

- (1) Primary productivity in the oceans is limited by the lack of nutrients in surface water.
- (2) These nutrients are mostly supplied from nutrient-rich subsurface waters through upwelling and vertical mixing (Barber 1992).
- (3) In ocean gyres upwelling and vertical mixing are not fully account for the observed productivity (Jenkins & Goldman 1985).
- (4) Upward pumping of nutrients is through action of meso-scale eddies (McGillicuddy et al. 1999, Segal et al. 1999).

Can we identify such an upward
pumping mechanism in western
Pacific (such as KE) using
satellite data?

Purpose of the Study

- To examine the spatial and temporal variability of SeaWiFS chlorophyll-*a* (Chl-*a*) in the KE region in relation to eddy structure
- To identify the upward nutrient pumping mechanism


Kuroshio Extension (KE)

- To the east of Japan the Kuroshio swings eastward to form the Kuroshio Extension. The branching of this current in the region of 160° E results in the movement known as the North Pacific Current.
- The Kuroshio Extension (KE) current carries warm water at nearly 140 million cubic meters per second (140 Sv) eastward into the North Pacific.

Sea Level Anomaly (SLA)

- SLA is measured by ERS 1/2 and TOPEX/Poseidon satellites at 7-day intervals.
- SLA has evident annual cycle and mesoscale structure dominated by eddy or Rossby waves.


Annual Sea Level amplitude (mm) in the North Pacific Ocean. The Kuroshio Extension route is associate with large annual amplitudes




Annual Signal of SLA

- (1) The annual signal is influenced partially by the ocean circulation and partially by the rise and fall of the sea surface that arises from the expansion and contraction of the ocean due to its heating and cooling with seasonal climatic change.
- (2) Maximum annual elevation change is about ± 20 cm

Topography of the studied area and Kuroshio Extension axis (marked orange) adopted for the present study. Stations positions 1, 15, 40 and 80 are marked.



Satellite observed sea level anomaly (cm) in the North Pacific in April 2007




Chl-*a* Concentration

- Chl-*a* concentration is computed from the ratio of radiances measured in band-3 (480–500 nm) and band-5 (545–565 nm) according to the following NASA algorithm,

-

$$C_a = \exp \left[0.464 - 1.989 \ln \left(L_{WM} 490 / L_{WM} 555 \right) \right]$$


Satellite observed Chl-a concentration (mg/m³) in the North Pacific in April 2007.



Chl-a concentration

Cyclonic eddy \leftrightarrow High Chl-a


Anticyclonic eddy \leftrightarrow Low Chl-a




SLA (mm)

Anticyclonic eddies \rightarrow
A, B, C, ...

Cyclonic eddies \rightarrow
1, 2, ...




Chl-*a* (mg m^{-3} Chl *a*) seasonal cycle along the route of the Kuroshio Extension. The results for station 1-15 are in grey and for station 16-80 in black




$$r = -0.45$$

**Correlation between
Chl-*a* and SLA residuals
along the route of the
Kuroshio Extension**



$$r = -0.43$$



Conclusions

- (1) Upward pumping of nutrients by meso-scale eddies in the KE region was identified.
- (3) SeaWiFS chlorophyll-*a* concentrations are redistributed at the eddy scale by the eddy surface swirl currents.
- (4) The zonal scale for chlorophyll perturbations was determined at ~ 460 km