



Contents lists available at ScienceDirect

Applied Mathematics and Computation

journal homepage: www.elsevier.com/locate/amc

Galerkin finite element method for one nonlinear integro-differential model

Temur Jangveladze^a, Zurab Kiguradze^b, Beny Neta^{c,*}

^a Ilia Vekua Institute of Applied Mathematics, Ivane Javakhishvili Tbilisi State University, 2 University St., 0186 Tbilisi, Georgia

^b Ivane Javakhishvili Tbilisi State University, 2 University St., 0186 Tbilisi, Georgia

^c Department of Applied Mathematics, Naval Postgraduate School Monterey, CA 93943, USA

ARTICLE INFO

Keywords:

Nonlinear integro-differential equations
Finite elements
Galerkin method

ABSTRACT

Galerkin finite element method for the approximation of a nonlinear integro-differential equation associated with the penetration of a magnetic field into a substance is studied. First type initial-boundary value problem is investigated. The convergence of the finite element scheme is proved. The rate of convergence is given too. The decay of the numerical solution is compared with the analytical results.

Published by Elsevier Inc.

1. Introduction

The goal of this paper is a study of Galerkin finite element method for approximation of a nonlinear integro-differential equation arising in mathematical modeling of the process of a magnetic field penetrating into a substance. If the coefficient of thermal heat capacity and electroconductivity of the substance highly dependent on temperature, then the Maxwell's system [1], that describe above-mentioned process, can be rewritten in the following form [2]:

$$\frac{\partial W}{\partial t} = -\text{rot} \left[a \left(\int_0^t |\text{rot} W|^2 d\tau \right) \text{rot} W \right], \quad (1.1)$$

where $W = (W_1, W_2, W_3)$ is a vector of the magnetic field and the function $a = a(\sigma)$ is defined for $\sigma \in [0, \infty)$. Let us consider magnetic field W , with the form $W = (0, 0, u)$, where $u = u(x, t)$ is a scalar function of time and of one spatial variables. Then $\text{rot} W = (0, -\frac{\partial u}{\partial x}, 0)$ and Eq. (1.1) will take the form

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial x} \left[a \left(\int_0^t \left(\frac{\partial u}{\partial x} \right)^2 d\tau \right) \frac{\partial u}{\partial x} \right]. \quad (1.2)$$

Note that (1.2) is complex, but special cases of such type models were investigated, see [2–11]. The existence of global solutions for initial-boundary value problems of such models have been proven in [2–4,10] by using the Galerkin and compactness methods [12,13]. The asymptotic behavior of the solutions of (1.2) have been the subject of intensive research in recent years (see e.g. [10,14–20]).

In [7] some generalization of equations of type (1.1) is proposed. There it was assumed that the temperature of the considered body is depending on time, but independent of the space coordinates. If the magnetic field again has the form $W = (0, 0, u)$ and $u = u(x, t)$, then the same process of penetration of the magnetic field into the material is modeled by the following integro-differential equation [7]:

* Corresponding author.

E-mail address: bneta@nps.edu (B. Neta).

$$\frac{\partial u}{\partial t} = a \left(\int_0^t \int_0^1 \left(\frac{\partial u}{\partial x} \right)^2 dx d\tau \right) \frac{\partial^2 u}{\partial x^2}. \tag{1.3}$$

The asymptotic behavior of solutions of the initial-boundary value problem for the Eq. (1.3) and the convergence of the finite difference approximation for the case $a(\sigma) = 1 + \sigma$ were studied in [17]. The solvability and uniqueness of the solutions of (1.3) type model is studied in [10].

Note that in [17,21–24] difference schemes for (1.2), (1.3) type models were investigated. Difference schemes for one nonlinear parabolic integro-differential model similar to (1.2) were studied in [25] and [26].

The purpose of this study is to develop a Galerkin finite element method to solve (1.3). The rest of the paper is organized as follows. In the next section the variational formulation of problem is given. In the third section finite element scheme for (1.3) is described and error estimate is proven. We close with a section on numerical implementation and present numerical results comparing with the theoretical ones.

2. Variational Formulation

Consider the following initial-boundary value problem:

$$\frac{\partial u}{\partial t} = (1 + \sigma(t)) \frac{\partial^2 u}{\partial x^2} + f(x, t), \quad (x, t) \in Q = (0, 1) \times (0, T), \tag{2.1}$$

$$u(0, t) = u(1, t) = 0, \quad t \geq 0, \tag{2.2}$$

$$u(x, 0) = u_0(x), \quad x \in [0, 1], \tag{2.3}$$

where

$$\sigma(t) = \int_0^t \int_0^1 \left(\frac{\partial u}{\partial x} \right)^2 dx d\tau$$

and $u_0(x)$ is a given function.

One of the ingredients of finite-element method is a variational formulation of the problem. Let us denote by H the linear space of functions u satisfying (2.2) and

$$\|u(\cdot, t)\|_1 < \infty,$$

where

$$\|u(\cdot, t)\|_r = \left\{ \int_0^1 \left[|u(x, t)|^2 + \sum_{i=1}^r \left| \frac{\partial^i u(x, t)}{\partial x^i} \right|^2 \right] dx \right\}^{1/2}.$$

The variational formulation of problem (2.1)–(2.3) can be stated as follows: Find a function $u(x, t) \in H$ for which

$$\left\langle v, \frac{\partial u}{\partial t} \right\rangle + \left\langle (1 + \sigma(t)) \frac{\partial u}{\partial x}, \frac{\partial v}{\partial x} \right\rangle = \langle f, v \rangle, \quad \forall v \in H, \tag{2.4}$$

and

$$\langle v, u(x, 0) \rangle = \langle v, u_0(x) \rangle, \quad \forall v \in H, \tag{2.5}$$

where $\langle p(x), q(x) \rangle = \int_0^1 p(x)q(x)dx$. To approximate the solution of (2.4), (2.5) we require that u lies in a finite-dimensional subspace S_h of H for each t (see e.g. [27]). The following property concerning approximability in S_h can be readily verified for finite-element spaces (see e.g. [28]).

2.1. Approximation Property

There is an integer $r \geq 2$ and numbers C_0, C_1 independent of h such that for any $v \in H$ there exists a $v^h \in S_h$ satisfying

$$\|v - v^h\|_\ell \leq C_\ell h^{r-\ell} \|v\|_j, \quad \text{for } 0 \leq \ell \leq 1, \quad \ell < j \leq r. \tag{2.6}$$

2.2. Approximate problem

The approximation $u^h \in S_h$ to u is defined by the following variational analog of (2.4), (2.5):

Find a $u^h \in S_h$ such that

$$\left\langle v^h, \frac{\partial u^h}{\partial t} \right\rangle + \left\langle (1 + \sigma_h(t)) \frac{\partial u^h}{\partial x}, \frac{\partial v^h}{\partial x} \right\rangle = \langle f, v^h \rangle, \quad \forall v^h \in S_h, \tag{2.7}$$

and

$$\langle v^h, u^h(x, 0) \rangle = \langle v^h, u_0(x) \rangle, \quad \forall v^h \in S_h, \tag{2.8}$$

where

$$\sigma_h(t) = \int_0^t \int_0^1 \left(\frac{\partial u^h}{\partial x} \right)^2 dx d\tau.$$

Once a basis has been selected for S_h , (2.7), (2.8) are equivalent to a set of N integro-differential equations, where N is dimension of S_h . The solution of such a system will be discussed in Section 4.

3. Error estimates

In this section we shall estimate the error in the finite element approximation using the norm

$$\| \| E \| \|_r = \int_0^T \int_0^1 \sum_{i=0}^r \left| \frac{\partial^i E(x, t)}{\partial x^i} \right|^2 dx dt.$$

Everywhere in the case $r = 0$ we will omitted the subscript.

Theorem. *The error in the finite element approximation u^h generated by (2.7), (2.8) satisfies the relation*

$$\| \| u - u^h \| \|_1 \leq h^{j-1} C \left\{ \frac{1}{2} h^2 \| u_0 \|^2 + Ch^2 \| \| u_t \| \|^2 + C^2 [1 + h^{2(j-1)} \| \| u \| \|^2] \| \| u \| \|^2 + C^3 (h^{j-1} \| \| u \| \|^2 + \| \| u \| \|^2) \right\}^{1/2},$$

where

$$\| \| u \| \| = \int_0^T \int_0^1 |u| dx d\tau.$$

Here and below C and c_i denote various constants independent of h .

Proof. Subtracting (2.7) from (2.4) with v^h instead of v we obtain

$$\left\langle v^h, \frac{\partial u^h}{\partial t} \right\rangle + \left\langle (1 + \sigma_h(t)) \frac{\partial u^h}{\partial x}, \frac{\partial v^h}{\partial x} \right\rangle = \left\langle v^h, \frac{\partial u}{\partial t} \right\rangle + \left\langle (1 + \sigma(t)) \frac{\partial u}{\partial x}, \frac{\partial v^h}{\partial x} \right\rangle, \quad \forall v^h \in S_h.$$

Let \tilde{u}^h be any function in S_h , then

$$\begin{aligned} & \left\langle v^h, \frac{\partial (u^h - \tilde{u}^h)}{\partial t} \right\rangle + \left\langle \left[(1 + \sigma_h(t)) \frac{\partial u^h}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial v^h}{\partial x} \right\rangle \\ & = \left\langle v^h, \frac{\partial (u - \tilde{u}^h)}{\partial t} \right\rangle + \left\langle \left[(1 + \sigma(t)) \frac{\partial u}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial v^h}{\partial x} \right\rangle, \quad \forall v^h \in S_h, \end{aligned} \tag{3.1}$$

where

$$\tilde{\sigma}_h(t) = \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau.$$

Let us define the errors as follows:

$$\begin{aligned} e(x, t) &= u^h(x, t) - \tilde{u}^h(x, t), \\ E(x, t) &= u(x, t) - \tilde{u}^h(x, t). \end{aligned} \tag{3.2}$$

Since $e \in S_h$, we can let $v^h = e$ and (3.1) becomes

$$\left\langle e, \frac{\partial e}{\partial t} \right\rangle + \left\langle \left[(1 + \sigma_h(t)) \frac{\partial u^h}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle = \left\langle e, \frac{\partial E}{\partial t} \right\rangle + \left\langle \left[(1 + \sigma(t)) \frac{\partial u}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle. \tag{3.3}$$

Let us consider the second term on the left of (3.3)

$$\left\langle \left[(1 + \sigma_h(t)) \frac{\partial u^h}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle = \left\langle \frac{\partial e}{\partial x}, \frac{\partial e}{\partial x} \right\rangle + \left\langle \left[\sigma_h(t) \frac{\partial u^h}{\partial x} - \tilde{\sigma}_h(t) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle. \tag{3.4}$$

Denoting

$$\omega = \frac{\partial u^h}{\partial x}, \quad \eta = \frac{\partial \tilde{u}^h}{\partial x},$$

the last term of (3.4) can be rewritten as

$$\begin{aligned} & \left\langle \left[\sigma_h(t) \frac{\partial u^h}{\partial x} - \tilde{\sigma}_h(t) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle \\ &= \left\langle \omega \int_0^t \int_0^1 \omega^2 dx d\tau - \eta \int_0^t \int_0^1 \eta^2 dx d\tau, \omega - \eta \right\rangle = \frac{1}{2} \left\langle \int_0^t \int_0^1 \omega^2 dx d\tau + \int_0^t \int_0^1 \eta^2 dx d\tau, (\omega - \eta)^2 \right\rangle \\ &+ \frac{1}{2} \left\langle \int_0^t \int_0^1 \omega^2 dx d\tau - \int_0^t \int_0^1 \eta^2 dx d\tau, \omega^2 - \eta^2 \right\rangle \geq \frac{1}{2} \left\langle \int_0^t \int_0^1 \omega^2 dx d\tau - \int_0^t \int_0^1 \eta^2 dx d\tau, \omega^2 - \eta^2 \right\rangle \\ &= \frac{1}{2} \int_0^t \int_0^1 (\omega^2 - \eta^2) dx d\tau (\omega^2 - \eta^2) dx = \frac{1}{2} \int_0^t \int_0^1 (\omega^2 - \eta^2) dx d\tau \int_0^1 (\omega^2 - \eta^2) dx = \frac{1}{4} \frac{d\varphi^2(t)}{dt}, \end{aligned}$$

where

$$\varphi(t) \equiv \int_0^t \int_0^1 (\omega^2 - \eta^2) dx d\tau.$$

Therefore, left hand side of (3.3) can be rewritten as follows

$$\left\langle e, \frac{\partial e}{\partial t} \right\rangle + \left\langle \left[(1 + \sigma_h(t)) \frac{\partial u^h}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle \geq \frac{1}{2} \frac{d}{dt} \|e\|^2 + \|e_x\|^2 + \frac{1}{4} \frac{d\varphi^2(t)}{dt}.$$

Now consider the second term on the right of (3.3)

$$\left\langle \left[(1 + \sigma(t)) \frac{\partial u}{\partial x} - (1 + \tilde{\sigma}_h(t)) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle = \left\langle \frac{\partial E}{\partial x}, \frac{\partial e}{\partial x} \right\rangle + \left\langle \left[\sigma(t) \frac{\partial u}{\partial x} - \tilde{\sigma}_h(t) \frac{\partial \tilde{u}^h}{\partial x} \right], \frac{\partial e}{\partial x} \right\rangle. \tag{3.5}$$

Substituting for u_x from (3.2), the last equality gives

$$\begin{aligned} \sigma(t) \frac{\partial u}{\partial x} - \tilde{\sigma}_h(t) \frac{\partial \tilde{u}^h}{\partial x} &= \left(E_x + \frac{\partial \tilde{u}^h}{\partial x} \right) \int_0^t \int_0^1 \left(E_x + \frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau - \frac{\partial \tilde{u}^h}{\partial x} \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau \\ &= \left(E_x + \frac{\partial \tilde{u}^h}{\partial x} \right) \int_0^t \int_0^1 \left(E_x^2 + 2E_x \frac{\partial \tilde{u}^h}{\partial x} \right) dx d\tau + E_x \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau. \end{aligned}$$

Taking this into account in the right hand side of (3.3) we get

$$\begin{aligned} & \langle e, E_t \rangle + \langle E_x, e_x \rangle + \langle E_x, e_x \rangle \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \langle E_x, e_x \rangle \int_0^t \int_0^1 E_x^2 dx d\tau + \left\langle \frac{\partial \tilde{u}^h}{\partial x}, e_x \right\rangle \int_0^t \int_0^1 E_x^2 dx d\tau + \left\langle E_x + \frac{\partial \tilde{u}^h}{\partial x}, e_x \right\rangle \\ & \times \int_0^t \int_0^1 2 \frac{\partial \tilde{u}^h}{\partial x} E_x dx d\tau = \langle e, E_t \rangle + \langle E_x, e_x \rangle \left\{ 1 + \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \int_0^t \int_0^1 E_x^2 dx d\tau + \int_0^t \int_0^1 2 \frac{\partial \tilde{u}^h}{\partial x} E_x dx d\tau \right\} \\ & + \left\langle \frac{\partial \tilde{u}^h}{\partial x}, e_x \right\rangle \left\{ \int_0^t \int_0^1 E_x^2 dx d\tau + \int_0^t \int_0^1 2 \frac{\partial \tilde{u}^h}{\partial x} E_x dx d\tau \right\} \leq \langle e, E_t \rangle + \langle |E_x|, |e_x| \rangle \left\{ 1 + \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau \right. \\ & + \int_0^t \int_0^1 E_x^2 dx d\tau + \frac{1}{\epsilon_1} \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \epsilon_1 \int_0^t \int_0^1 E_x^2 dx d\tau \left. \right\} + \left\langle \left| \frac{\partial \tilde{u}^h}{\partial x} \right|, |e_x| \right\rangle \left\{ \int_0^t \int_0^1 E_x^2 dx d\tau \right. \\ & + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \int_0^t \int_0^1 |E_x| dx d\tau \left. \right\} \leq \langle e, E_t \rangle + \langle |E_x|, |e_x| \rangle \left\{ 1 + \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau \right. \\ & + \int_0^t \int_0^1 E_x^2 dx d\tau + \frac{1}{\epsilon_1} \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \epsilon_1 \int_0^t \int_0^1 E_x^2 dx d\tau \left. \right\} + \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \int_0^1 |e_x| dx \left\{ \int_0^t \int_0^1 E_x^2 dx d\tau \right. \\ & + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \int_0^t \int_0^1 |E_x| dx d\tau \left. \right\}, \end{aligned}$$

where $\epsilon_1 > 0$ comes from Schwartz inequality.

Now incorporate these estimates into (3.3) to have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|e\|^2 + \|e_x\|^2 + \frac{1}{4} \frac{d\varphi^2(t)}{dt} \leq \langle e, E_t \rangle + \langle |E_x|, |e_x| \rangle \left\{ 1 + \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \int_0^t \int_0^1 E_x^2 dx d\tau \right. \\ & + \frac{1}{\epsilon_1} \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x} \right)^2 dx d\tau + \epsilon_1 \int_0^t \int_0^1 E_x^2 dx d\tau \left. \right\} + \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \int_0^1 |e_x| dx \left\{ \int_0^t \int_0^1 E_x^2 dx d\tau \right. \\ & + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \int_0^t \int_0^1 |E_x| dx d\tau \left. \right\}. \end{aligned}$$

Integrate with respect to t , we have

$$\begin{aligned} & \frac{1}{2} \|e(\cdot, T)\|^2 + \int_0^T \|e_x\|^2 dt + \varphi^2(T) + \leq \frac{1}{2} \|e(\cdot, 0)\|^2 + \int_0^T \int_0^1 e E_t dx dt \\ & + \int_0^T \int_0^1 |E_x e_x| dx \left\{ 1 + (1 + \epsilon_1) \int_0^t \int_0^1 E_x^2 dx d\tau + \left(1 + \frac{1}{\epsilon_1}\right) \int_0^t \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x}\right)^2 dx d\tau \right\} dt \\ & + \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \int_0^1 |e_x| dx \left\{ \int_0^t \int_0^1 E_x^2 dx d\tau + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \int_0^t \int_0^1 |E_x| dx d\tau \right\} dt. \end{aligned} \tag{3.6}$$

Note that the first, third and fourth terms on the left are nonnegative and can be dropped. We also use Schwarz inequality on the right hand side for these two terms

$$\begin{aligned} \int_0^T \int_0^1 e E_t dx dt & \leq \frac{1}{2} (\epsilon_2 \| \|e\| \|^2 + 1/\epsilon_2 \| \|E_t\| \|^2), \\ \int_0^T \int_0^1 |E_x e_x| dx & \leq \frac{1}{2} (\epsilon_3 \| \|E_x\| \|^2 + 1/\epsilon_3 \| \|e_x\| \|^2). \end{aligned}$$

We can estimate the last term on the right by estimating the term in parenthesis and then take it outside the time integral

$$\begin{aligned} I & \equiv \int_0^t \int_0^1 E_x^2 dx d\tau + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \int_0^t \int_0^1 |E_x| dx d\tau \leq \| \|E_x\| \|^2 + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \int_0^T \int_0^1 |E_x| dx d\tau \\ & = \| \|E_x\| \|^2 + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \| \|E_x\| \|. \end{aligned}$$

Therefore, the last term in (3.6) becomes

$$\begin{aligned} I \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \int_0^1 |e_x| dx dt & \leq I \sqrt{\int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt} \sqrt{\int_0^T \left(\int_0^1 |e_x| dx \right)^2 dt} \leq I \sqrt{\int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt} \| \|e_x\| \|^2 \\ & \leq \frac{\epsilon_4}{2} I^2 \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt + \frac{1}{2\epsilon_4} \| \|e_x\| \|^2. \end{aligned}$$

Therefore, (3.6) gives

$$\begin{aligned} \| \|e_x\| \|^2 & \leq \frac{1}{2} \| \|e(\cdot, 0)\| \|^2 + \frac{\epsilon_2}{2} \| \|e\| \|^2 + \frac{1}{2\epsilon_2} \| \|E_t\| \|^2 + M_1 \left(\frac{\epsilon_3}{2} \| \|E_x\| \|^2 + \frac{1}{2\epsilon_3} \| \|e_x\| \|^2 \right) + (1 + \epsilon_1) \| \|E_x\| \|^2 \left(\frac{\epsilon_3}{2} \| \|E_x\| \|^2 + \frac{1}{2\epsilon_3} \| \|e_x\| \|^2 \right) \\ & + \frac{\epsilon_4}{2} I^2 \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt + \frac{1}{2\epsilon_4} \| \|e_x\| \|^2, \end{aligned}$$

where

$$M_1 \equiv 1 + \left(1 + \frac{1}{\epsilon_1}\right) \int_0^T \int_0^1 \left(\frac{\partial \tilde{u}^h}{\partial x}\right)^2 dx d\tau.$$

Combining terms we have

$$\begin{aligned} \left[1 - \frac{M_1}{2\epsilon_3} - \frac{1}{2\epsilon_4} - \frac{1 + \epsilon_1}{2\epsilon_3} \| \|E_x\| \|^2 \right] \| \|e_x\| \|^2 - \frac{\epsilon_2}{2} \| \|e\| \|^2 & \leq \frac{1}{2} \| \|e(\cdot, 0)\| \|^2 + \frac{1}{2\epsilon_2} \| \|E_t\| \|^2 + \left[\frac{M_1 \epsilon_3}{2} + \frac{(1 + \epsilon_1) \epsilon_3}{2} \| \|E_x\| \|^2 \right] \| \|E_x\| \|^2 \\ & + \frac{\epsilon_4}{2} \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt \left(\| \|E_x\| \|^2 + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \| \|E_x\| \|^2 \right). \end{aligned}$$

Now we use Poincare' inequality

$$\| \|e_x\| \geq C_p \| \|e\| \|$$

to show that

$$\| \|e_x\| \geq C_p \| \|e\| \|$$

for possibly different constant C_p .

We have

$$\begin{aligned} \left[1 - \frac{M_1}{2\epsilon_3} - \frac{1}{2\epsilon_4} - \frac{1 + \epsilon_1}{2\epsilon_3} \| \|E_x\| \|^2 - \frac{\epsilon_2}{2C_p^2} \right] \| \|e_x\| \|^2 & \leq \frac{1}{2} \| \|e(\cdot, 0)\| \|^2 + \frac{1}{2\epsilon_2} \| \|E_t\| \|^2 + \left[\frac{M_1 \epsilon_3}{2} + \frac{(1 + \epsilon_1) \epsilon_3}{2} \| \|E_x\| \|^2 \right] \| \|E_x\| \|^2 \\ & + \frac{\epsilon_4}{2} \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right|^2 dt \left(\| \|E_x\| \|^2 + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x}\right) \right| \| \|E_x\| \|^2 \right). \end{aligned} \tag{3.7}$$

Note that we can choose $\epsilon_i, i = 1, \dots, 4$ so that the coefficient of the term on the left hand side of (3.7)

$$M_2 \equiv 1 - \frac{M_1}{2\epsilon_3} - \frac{1}{2\epsilon_4} - \frac{1 + \epsilon_1}{2\epsilon_3} \|E_x\|^2 - \frac{\epsilon_2}{2C_p^2}$$

is a positive constant. Recall that

$$\|e\|_1^2 = \|e\|^2 + \|e_x\|^2 \leq \left(1 + \frac{1}{C_p^2}\right) \|e_x\|^2,$$

so using (3.7) we get

$$\|e\|_1^2 \leq c_1^2 \left\{ \frac{1}{2} \|e(\cdot, 0)\|^2 + \frac{1}{2\epsilon_2} \|E_t\|^2 + \left[\frac{M_1\epsilon_3}{2} + \frac{(1 + \epsilon_1)\epsilon_3}{2} \|E_x\|^2 \right] \|E_x\|^2 + c_2 \left(\|E_x\|^2 + 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| \|E_x\| \right)^2 \right\},$$

where

$$c_1 = \sqrt{\frac{\left(1 + \frac{1}{C_p^2}\right)}{M_2}}$$

and

$$c_2 = \frac{\epsilon_4}{2} \int_0^T \left| \sup_x \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right|^2 dt.$$

Since E is the interpolation error, from (2.6) we have

$$\begin{aligned} \|E\| &\leq C_0 h^j \|u\|, \\ \|E_x\| &\leq C_1 h^{j-1} \|u\|, \\ \|E_t\| &\leq C_2 h^j \|u_t\|, \\ \|E_x\| &\leq C_3 h^{j-1} \|u\|, \end{aligned}$$

which yield

$$\|e\|_1 \leq h^{j-1} c_1 \left\{ \frac{1}{2} h^2 \|u_0\|^2 + c_3 h^2 \|u_t\|^2 + [c_4 + c_5 h^{2(j-1)} \|u\|^2] c_7 \|u\|^2 + c_2 (c_7 h^{j-1} \|u\|^2 + c_6 \|u\|)^2 \right\}^{1/2}, \tag{3.8}$$

where

$$c_3 = \frac{1}{2\epsilon_2} C_2^2, \quad c_4 = \frac{M_1\epsilon_3}{2}, \quad c_5 = \frac{(1 + \epsilon_1)\epsilon_3}{2} C_1^2, \quad c_6 = 2 \left| \sup_{x,t} \left(\frac{\partial \tilde{u}^h}{\partial x} \right) \right| C_3, \quad c_7 = C_1^2. \tag{3.9}$$

Using the triangle inequality we get

$$\|u - u^h\|_1^2 = \|u - \bar{u}^h + \bar{u}^h - u^h\|_1^2 \leq \|E\|_1^2 + \|e\|_1^2.$$

Therefore,

$$\|u - u^h\|_1^2 \leq c_8^2 h^{2(j-1)} \|u\|^2 + h^{2(j-1)} c_1^2 \left\{ \frac{1}{2} h^2 \|u_0\|^2 + c_3 h^2 \|u_t\|^2 + [c_4 + c_5 h^{2(j-1)} \|u\|^2] c_7 \|u\|^2 + c_2 (c_7 h^{j-1} \|u\|^2 + c_6 \|u\|)^2 \right\},$$

or

$$\|u - u^h\|_1^2 \leq h^{2(j-1)} c_1^2 \left\{ \frac{1}{2} h^2 \|u_0\|^2 + c_3 h^2 \|u_t\|^2 + [c_9 + c_5 h^{2(j-1)} \|u\|^2] c_7 \|u\|^2 + c_2 (c_7 h^{j-1} \|u\|^2 + c_6 \|u\|)^2 \right\}, \tag{3.10}$$

where $c_9 = c_4 + \frac{c_5^2}{c_1^2 c_7}$. From (3.10), by denoting $C = \max_{i=1,2,\dots,9} \{c_i\}$ we complete proof of theorem.

4. Numerical Solution

For the numerical solution of (2.7), (2.8) we let $\phi_1(x), \dots, \phi_N(x)$ be a basis for S_h . Therefore $u^h \in S_h$ can be represented by

$$u^h(x, t) = \sum_{j=1}^N u_j(t) \phi_j(x). \tag{4.1}$$

Since (2.7), (2.8) are valid for all $v^h \in S_h$, one can let $v^h = \phi_k$. This yields the following system for the weights $\mathbf{u}(t)$:

$$M\dot{\mathbf{u}} + K(u)\mathbf{u} = \mathbf{F}, \tag{4.2}$$

$$M\mathbf{u}(0) = \mathbf{U}, \tag{4.3}$$

where

$$M_{jk} = \langle \phi_k, \phi_j \rangle, \tag{4.4}$$

$$K(u)_{jk} = \langle (1 + \sigma_h(t))\phi'_k, \phi'_j \rangle, \tag{4.5}$$

$$\mathbf{F}_j = \langle \phi_j, f \rangle, \quad \mathbf{U}_j = \langle \phi_j, u_0 \rangle. \tag{4.6}$$

Now we can evaluate $\sigma_h(t)$ as follows

$$\begin{aligned} \sigma_h(t) &= \int_0^t \int_0^1 \left(\sum_{\ell=1}^N u_\ell(\tau)\phi'_\ell(x) \right)^2 dx d\tau = \int_0^t \int_0^1 \sum_{\ell=1}^N \sum_{m=1}^N u_\ell(\tau)u_m(\tau)\phi'_\ell(x)\phi'_m(x) dx d\tau \\ &= \sum_{\ell=1}^N \sum_{m=1}^N \int_0^t u_\ell(\tau)u_m(\tau) \underbrace{\left[\int_0^1 \phi'_\ell(x)\phi'_m(x) dx \right]}_{=K_{\ell m}} d\tau = \sum_{\ell=1}^N \sum_{m=1}^N \tilde{K}_{\ell m} \int_0^t u_\ell(\tau)u_m(\tau) d\tau. \end{aligned} \tag{4.7}$$

The time integral can be approximated by the trapezoidal rule ($0 = t_0 < t_1 < t_2 < \dots < t_n = t$) as follows

$$\int_0^t u_\ell(\tau)u_m(\tau) d\tau = \sum_{p=0}^n \Delta t \zeta_p u_\ell(t_p)u_m(t_p), \tag{4.8}$$

where $\zeta_p = 1/2$ for $p = 0, n$ and $\zeta_p = 1$ for $p = 1, \dots, n - 1$. Combining (4.8) and (4.7) with (4.5), we get

$$\begin{aligned} K(u)_{jk} &= \left\langle \left(1 + \sum_{\ell=1}^N \sum_{m=1}^N \tilde{K}_{\ell m} \sum_{p=0}^n \Delta t \zeta_p u_\ell(t_p)u_m(t_p) \right) \phi'_k, \phi'_j \right\rangle = \left(1 + \sum_{\ell=1}^N \sum_{m=1}^N \tilde{K}_{\ell m} \sum_{p=0}^n \Delta t \zeta_p u_\ell(t_p)u_m(t_p) \right) \tilde{K}_{jk} \\ &= \left(1 + \sum_{p=0}^n \Delta t \zeta_p v(t_p) \right) \tilde{K}_{jk}, \end{aligned} \tag{4.9}$$

where $v(t) = \mathbf{u}^T(t)\tilde{K}\mathbf{u}(t)$. To solve the system (4.2) and (4.3), we use Taylor's series. Let

$$\mathbf{u}(t + \Delta t) = \mathbf{u}(t) + (\Delta t)\dot{\mathbf{u}}(t) + \frac{1}{2}(\Delta t)^2\ddot{\mathbf{u}}(t) + O((\Delta t)^3). \tag{4.10}$$

Differentiating (4.2) with respect to t , one has

$$M\ddot{\mathbf{u}} + K(u)\dot{\mathbf{u}} + \dot{K}\mathbf{u} = \dot{\mathbf{F}}, \tag{4.11}$$

where

$$\dot{K}_{kj} = \langle \dot{\sigma}_h \phi'_j, \phi'_k \rangle = \left\langle \int_0^1 \left(\sum_{\ell=1}^N u_\ell(t)\phi'_\ell \right)^2 dx \phi'_j, \phi'_k \right\rangle = \sum_{\ell=1}^N \sum_{m=1}^N u_\ell(t)u_m(t)\tilde{K}_{\ell m}\tilde{K}_{kj} = \mathbf{u}^T(t)\tilde{K}\mathbf{u}(t)\tilde{K}_{kj} = v(t)\tilde{K}_{kj}. \tag{4.12}$$

Now multiply (4.10) by M and using (4.2), (4.11) and (4.12), we have after dropping terms of order higher than second

$$\begin{aligned} M(\mathbf{u}(t + \Delta t) - \mathbf{u}(t)) &= (\Delta t)M\dot{\mathbf{u}}(t) + \frac{1}{2}(\Delta t)^2M\ddot{\mathbf{u}}(t) = (\Delta t)[\mathbf{F} - K(u)\mathbf{u}] + \frac{1}{2}(\Delta t)^2[\dot{\mathbf{F}} - K(u)\dot{\mathbf{u}} - \dot{K}\mathbf{u}] \\ &= (\Delta t)[\mathbf{F} - K(u)\mathbf{u}] + \frac{1}{2}(\Delta t)^2[\dot{\mathbf{F}} - K(u)M^{-1}(\mathbf{F} - K(u)\mathbf{u}) - \dot{K}\mathbf{u}] \\ &= (\Delta t)\left[\mathbf{F} + \frac{1}{2}(\Delta t)\dot{\mathbf{F}} - \frac{1}{2}(\Delta t)K(u)M^{-1}\mathbf{F} \right] - (\Delta t)K(u)\left[\mathbf{u} - \frac{1}{2}\Delta tM^{-1}K(u)\mathbf{u} \right] - \frac{1}{2}(\Delta t)^2\dot{K}\mathbf{u}. \end{aligned} \tag{4.13}$$

If we take $t = t_n$ as in (4.8) and we denote $\mathbf{u}^n = \mathbf{u}(t_n)$ then substituting for K and \dot{K} from (4.9) and (4.12), we get

$$\begin{aligned} M(\mathbf{u}^{n+1} - \mathbf{u}^n) &= (\Delta t)\left[\mathbf{F}^n + \frac{1}{2}(\Delta t)\dot{\mathbf{F}}^n - \frac{1}{2}(\Delta t)\left(1 + \sum_{p=0}^n \Delta t \zeta_p v(t_p) \right) \tilde{K}M^{-1}\mathbf{F}^n \right] \\ &\quad - (\Delta t)\left(1 + \sum_{p=0}^n \Delta t \zeta_p v(t_p) \right) \tilde{K}\left[\mathbf{u}^n - \frac{1}{2}\Delta tM^{-1}\left(1 + \sum_{p=0}^n \Delta t \zeta_p v(t_p) \right) \tilde{K}\mathbf{u}^n \right] - \frac{1}{2}(\Delta t)^2 v(t)\tilde{K}\mathbf{u}^n. \end{aligned} \tag{4.14}$$




Fig. 1. The solution at $t = 0.5$. The exact solution is solid line and the numerical solution is marked by *.




Fig. 2. The solution at $t = 1.0$. The exact solution is solid line and the numerical solution is marked by *.

Table 1

The errors for various values of the grid spacing and the approximate rate of convergence.

h	Error	Rate
.2	.0071411	.964499
.04	.0015122	.978861
.02	.00076726	.988986
.01	.00038657	.994375
.005	.00019404	

Now let us denote

$$\psi^n = 1 + \sum_{p=0}^n \Delta t \zeta_p \nu^p, \tag{4.15}$$

then

$$M\mathbf{u}^{n+1} = M\mathbf{u}^n - (\Delta t)\psi^n \tilde{K} \left[\mathbf{u}^n - \frac{1}{2} \Delta t M^{-1} \psi^n \tilde{K} \mathbf{u}^n \right] - \frac{1}{2} (\Delta t)^2 \nu^n \tilde{K} \mathbf{u}^n + (\Delta t) \left[\mathbf{F}^n + \frac{1}{2} (\Delta t) \dot{\mathbf{F}}^n - \frac{1}{2} (\Delta t) \psi^n \tilde{K} M^{-1} \mathbf{F}^n \right]. \tag{4.16}$$

In our first numerical experiment we have chosen the right hand side so that the exact solution is given by

$$u(x, t) = x(1 - x) \sin(x + t).$$

In this case the right hand side is

$$f(x, t) = x(1 - x) \cos(x + t) - \left(1 + \frac{11}{60}t - \frac{1}{8} \cos(t) \sin(t) - \frac{1}{8} \cos(1 + t) \sin(1 + t) + \frac{1}{8} \cos(1) \sin(1) \right) \times (-2 \sin(x + t) + 2(1 - x) \cos(x + t) - 2x \cos(x + t) - x(1 - x) \sin(x + t)).$$

The parameters used are $M = 100$ which dictates $h = 0.01$. In the next two figures we plotted the numerical solution (marked with *) and the exact solution at $t = 0.5$ (Fig. 1) and $t = 1.0$ (Fig. 2) and it is clear that the two solutions are almost identical.

We have ran the same example with various values of h and measured the error $\|u - u_h\|_1$. The results are given in Table 1.

We have experimented with several other initial solutions, and in all cases we noticed the agreement with the exact solution.

Remark. Clearly, in general, the finite element approximation requires larger storage. In this case, the nonlinear system was solved using Taylor series approximation and one has to solve a banded system at each time step. No iteration is required. On the other hand, in the finite difference case, we have solved the nonlinear system using Newton’s method which required iterating with dense matrices. Both methods yield comparable numerical results.

References

[1] L. Landau, E. Lifschitz, *Electrodynamics of Continuous Media. Course of Theoretical Physics, vol. 8.* (Translated from the Russian) Pergamon Press, Oxford-London-New York-Paris; Addison-Wesley Publishing Co., Inc., Reading, Mass., 1960; Russian original: Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1957.

[2] D. Gordeziani, T. Dzhangveladze(Jangveladze), T. Korshia, Existence and uniqueness of the solution of a class of nonlinear parabolic problems, *Differ. Uravn.* 19 (1983) 1197–1207 (Russian). English translation: *Differ. Equat.* 19 (1983), 887–895.

[3] T. Dzhangveladze(Jangveladze), The first boundary value problem for a nonlinear equation of parabolic type, *Dokl. Akad. Nauk SSSR* 269 (1983) 839–842 (Russian). English translation: *Soviet Phys. Dokl.*, 28, (1983), 323–324.

[4] T. Dzhangveladze (Jangveladze), *An Investigation of the First Boundary-Value Problem for Some Nonlinear Parabolic Integrodifferential Equations*, Tbilisi State University, 1983. in Russian.

[5] G. Laptev, Quasilinear parabolic equations which contains in coefficients Volterra’s operator, *Math. Sbornik* 136 (1988) 530–545 (Russian). English translation: *Sbornik Math.* 64 (1989), 527–542.

[6] G. Laptev, Mathematical singularities of a problem on the penetration of a magnetic field into a substance, *Zh. Vychisl. Mat. Mat. Fiz.* 28 (1988) 1332–1345 (Russian). English translation: *U.S.S.R. Comput. Math. Math. Phys.*, 28, (1990), 35–45.

[7] G. Laptev, *Quasilinear Evolution Partial Differential Equations with Operator Coefficients*, Doctoral Dissertation, Moscow, 1990, (Russian).

[8] Y. Lin, H.M. Yin, Nonlinear parabolic equations with nonlinear functionals, *J. Math. Anal. Appl.* 168 (1992) 28–41.

[9] N. Long, A. Dinh, Nonlinear parabolic problem associated with the penetration of a magnetic field into a substance, *Math. Mech. Appl. Sci.* 16 (1993) 281–295.

[10] T.A. Jangveladze, On one class of nonlinear integro-differential equations, *Semin. I. Vekua Inst. Appl. Math.* 23 (1997) 51–87.

[11] Y. Bai, P. Zhang, On a class of Volterra nonlinear equations of parabolic type, *Appl. Math. Comput.* 216 (2010) 236–240.

[12] N. Vishik, Solvability of boundary-value problems for quasi-linear parabolic equations of higher orders (Russian), *Math. Sb. (N.S.)* 59 (101)(1962) 289–325. suppl.

[13] J. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non-linéaires*, Dunod, Gauthier-Villars, Paris, 1969.

[14] T. Jangveladze(Dzhangveladze), Z. Kiguradze, Asymptotics of a solution of a nonlinear system of diffusion of a magnetic field into a substance, *Sibirsk. Mat. Zh.* 47 (2006) 1058–1070 (Russian). English translation: *Siberian Math. J.*, 47, (2006), 867–878.

- [15] T. Jangveladze(Dzhangveladze), Z. Kiguradze, On the stabilization of solutions of an initial-boundary value problem for a nonlinear integro-differential equation, *Differ. Uravn.* 43 (2007) 833–840 (Russian). English translation: *Diff. Eq.*, 43, (2007), 854–861.
- [16] T. Jangveladze(Dzhangveladze), Z. Kiguradze, Asymptotic behavior of the solution to nonlinear integro-differential diffusion equation, *Differ. Uravn.* 44 (2008) 517–529 (Russian). English translation: *Diff. Eq.*, 44, (2008), 538–550.
- [17] T.A. Jangveladze, Z.V. Kiguradze, B. Neta, Large time behavior of solutions and finite difference scheme to a nonlinear integro-differential equation, *Comput. Math. Appl.* 57 (2009) 799–811.
- [18] T. Jangveladze, Z. Kiguradze, B. Neta, Large time behavior of solutions to a nonlinear integro-differential system, *J. Math. Anal. Appl.* 351 (2009) 382–391.
- [19] M. Aptsiauri, T. Jangveladze, Z. Kiguradze, On asymptotic behavior of solution of one nonlinear one-dimensional integro-differential analogue of Maxwell's system, *Rep. Enl. Sess. Sem. I.Vekua Inst. Appl. Math.* 23 (2009) 3–6.
- [20] T. Jangveladze, Z. Kiguradze, B. Neta, Large time asymptotic and numerical solution of a nonlinear diffusion model with memory, *Comput. Math. Appl.* 59 (2010) 254–273.
- [21] T.A. Jangveladze, Convergence of a difference scheme for a nonlinear integro-differential equation, *Proc. I. Vekua Inst. Appl. Math.* 48 (1998) 38–43.
- [22] Z. Kiguradze, The difference scheme for one system of nonlinear partial differential equations, *Rep. Enl. Sess. Sem. I.Vekua Inst. Appl. Math.* 14 (1999) 67–70.
- [23] Z.V. Kiguradze, Finite difference scheme for a nonlinear integro-differential system, *Proc. I. Vekua Inst. Appl. Math.* 50–51 (2000-2001) 65–72.
- [24] T. Jangveladze, Z. Kiguradze, B. Neta, Finite difference approximation of a nonlinear integro-differential system, *Appl. Math. Comput.* 215 (2009) 615–628.
- [25] B. Neta, Numerical solution of a nonlinear integro-differential equation, *J. Math. Anal. Appl.* 89 (1982) 598–611.
- [26] B. Neta, J.O. Igwe, Finite differences versus finite elements for solving nonlinear integro-differential equations, *J. Math. Anal. Appl.* 112 (1985) 607–618.
- [27] B. Neta, Finite element approximation of a nonlinear parabolic problem, *Comput. Math. Appl.* 4 (1978) 247–255.
- [28] G. Strang, G.J. Fix, *An Analysis of the Finite Element Methods*, Prentice-Hall, Englewood Cliffs, NJ, 1973.