

**Results:** There was a significant interaction between time and treatment for bodyweight gain ( $F_{(8,144)}=2.7$ ,  $P < 0.0082$ ) and food intake ( $F_{(8,144)}=6.1$ ,  $P < 0.0001$ ). Noise-exposed females gained significantly more weight and had greater food intake after 17-days of noise exposure compared to controls (weight gain (g):  $29.8 \pm 4.7$  vs.  $12.8 \pm 3.0$ ,  $P < 0.0001$ . food intake (g):  $367.2 \pm 5.0$  vs.  $335.3 \pm 7.6$ ,  $P < 0.0001$ , respectively). Weight gain and food intake was significantly greater in noise-exposed females on all treatment days ( $P < 0.05$ ) excluding treatment day 5 for weight gain and treatment days 3 and 5 for food intake.

**Conclusion:** The increase in weight gain and feeding due to noise-induced sleep disruption is similar in male and female rats. These data have implications for future therapies to combat disordered sleep and obesity among women and identify mechanisms underlying sex-dependent differences in sleep disparities due to noise and obesity.

**Support (If Any):** Department of Veterans Affairs, USDA, University of AZ.

## 0169

### SLEEP PATTERNS DURING DUTY PERIODS AND DURING OFF-DAYS BETWEEN DUTY CYCLES IN HOSPITAL EMPLOYEES WORKING 12-HOUR NIGHT SHIFTS

Riedy SM<sup>1</sup>, English A<sup>2</sup>, Albritton S<sup>2</sup>, Burton J<sup>2</sup>, Himmel M<sup>2</sup>, Morgan P<sup>2</sup>, Kadel KL<sup>1</sup>, Van Dongen H<sup>1</sup>, Wilson M<sup>3</sup>

<sup>1</sup>Sleep and Performance Research Center, Washington State University, Spokane, WA, <sup>2</sup>Kootenai Health, Coeur d'Alene, ID, <sup>3</sup>College of Nursing, Washington State University, Spokane, WA

**Introduction:** Around-the-clock operations are common in hospital settings. Employees working nights are forced by their work schedules to work at times when their circadian clock promotes sleep, and sleep at times when the circadian clock promotes wakefulness. Little is known about hospital night workers' sleep patterns during days off when not constrained by work schedules. We compared the sleep patterns of 12-hour night and day workers at two regional hospitals during duty cycles and during intervening off-days.

**Methods:** An online survey was accessible for one month to the employees of two regional hospitals in the USA's Inland Northwest. 1,340 surveys were submitted. Following data quality control, responses of 12-hour shift workers with duty start times of either 07:00 or 19:00 were selected, yielding  $n=193$  day worker surveys and  $n=134$  night worker surveys. We investigated reported bedtimes during duty cycles and between-cycles; global scores on the Pittsburgh Sleep Quality Index (PSQI); and subjective sleepiness on the Epworth Sleepiness Scale (ESS).

**Results:** Night workers went to bed in the morning during duty cycles (mean  $\pm$  SD:  $08:53 \pm 46$ min) and in the evening during off-days ( $23:53 \pm 152$ min), whereas day workers maintained bedtimes in the morning during both duty cycles ( $22:00 \pm 53$ min) and off-days ( $22:40 \pm 63$ min) (mixed-effects ANOVA interaction:  $F=112.6$ ,  $P < 0.001$ ). Compared to day workers, night workers exhibited greater PSQI scores (day: 7.1, night: 8.3; Wilcoxon signed-rank test:  $Z=2.89$ ,  $P=0.004$ ) and ESS scores (day: 6.9, night: 8.0;  $Z=2.18$ ,  $P=0.030$ ). PSQI scores were  $>5$  (clinically relevant poor sleep quality) for both groups.

**Conclusion:** Among hospital employees working 12-hour shifts, night workers reported significantly poorer sleep quality and greater subjective sleepiness than day workers. Self-reported bedtimes revealed that during off-days between duty cycles, the vast majority of night workers reverted to a nighttime sleep pattern similar to that of day workers.

**Support (If Any):** Pullman Regional Hospital and Kootenai Health.

## 0170

### SLEEPINESS AND DRIVING INCIDENTS IN NURSES COMMUTING TO AND FROM WORK SHIFTS

Mulhall MD<sup>1,2</sup>, Sletten TL<sup>1,2</sup>, Magee M<sup>1,2</sup>, Howard ME<sup>3,1,2</sup>, Rajaratnam SM<sup>1,2</sup>

<sup>1</sup>Monash Institute of Cognitive and Clinical Neurosciences, School of Psychology, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, AUSTRALIA, <sup>2</sup>Cooperative Research Centre for Alertness, Safety and Productivity, Melbourne, AUSTRALIA, <sup>3</sup>Institute for Breathing and Sleep, Austin Health, Melbourne, AUSTRALIA

**Introduction:** Shift workers are at increased risk of sleepiness and motor vehicle accidents. This study aimed to assess sleepiness and adverse driving events during commutes to and from work shifts in intensive care unit (ICU) nurses.

**Methods:** 35 ICU nurses (age  $33.2 \pm 10.6$  years, 28 Female) completed driving diaries for commutes to and from day (07:00-15:30h), evening (13:00-21:00h), and night shifts (21:00-07:00h), consisting of a Karolinska Sleepiness Scale (KSS) pre- and post-drive and driving incidents. Optalert<sup>TM</sup> was used to monitor drowsiness during all drives. Linear mixed model analyses were conducted to investigate differences in pre-drive and post-drive KSS and driving incidents between shift types, controlling for number of drives per participant and drive duration.

**Results:** On the commute home from work, night shift was associated with the highest pre-drive KSS ( $6.31 \pm 1.73$ ), compared to evenings ( $3.92 \pm 1.60$ ) and days ( $3.80 \pm 1.76$ ,  $p < 0.001$ ). Significantly more driving incidents were reported on the commute home from night shift ( $3.01 \pm 2.09$ ) than on days ( $2.00 \pm 1.53$ ) or evenings ( $1.76 \pm 1.27$ ,  $p < 0.01$ ). In contrast, on commutes to work, pre-drive KSS scores were highest on day shift ( $5.06 \pm 1.79$ ), followed by nights ( $4.10 \pm 1.60$ ) and evenings ( $2.61 \pm 1.36$ ,  $p < 0.001$ ). Post-drive KSS was significantly higher on the drive to day ( $4.63 \pm 1.67$ ) and night shifts ( $4.78 \pm 1.53$ ) compared to evening shifts ( $2.72 \pm 1.42$ ,  $p < 0.001$ ). The most frequently reported driving incidents were: being distracted (13%), fighting to stay awake (10%), and fixation on interior/exterior objects (10%).

**Conclusion:** Nurses driving home from night shift are at increased risk of sleepiness and adverse driving events. Increased sleepiness was also observed during commute to day shifts compared to other shift types, likely due to the combined effects of the circadian drive for sleepiness and sleep restriction due to evening- to day-shift rotations (reported in 74% of participants). Identification of high-risk commutes may assist in the development of sleepiness monitoring and intervention strategies.

**Support (If Any):** The study was supported by the Cooperative Research Centre for Alertness, Safety and Productivity.

## 0171

### WORK AND SLEEP PATTERNS IN MILITARY SHIFT WORKERS: PROMOTING HEALTH AND WELLNESS THROUGH INFORMED SHIFT SCHEDULES

Shattuck NL, Matsangas P

Naval Postgraduate School, Monterey, CA

**Introduction:** Reduced manning, extended work hours, and circadian misalignment are some of the factors which contribute to the sleep debt and degraded alertness observed in much of the military population. This study compared fatigue levels, sleep, and real time performance measures (e.g., reaction time) of military shift workers.

**Methods:** Working either 8-hour (daytime, evening, or overnight) or 12-hour (day, overnight) shifts, military shift workers ( $N=75$ ) serving

in various locations around the world participated in the 2-week study. Participants wore actigraphs, completed activity logs, and performed a three-minute Psychomotor Vigilance Task (PVT) before and after standing watch. In addition, they completed sleep habits and mood states questionnaires at the beginning and end of the study.

**Results:** Participants slept 6.74 hours/day, with 19% sleeping less than 6 hours on average. At the outset of the study, ~62% of the participants reported insomnia symptoms and were classified as poor sleepers. Although sleep duration did not differ between schedules, participants on the 8-hour shifts made fewer errors and showed less variability in PVT (e.g., fewer lapses combined with false starts) compared to those individuals working on 12-hour shifts. Participants on 12-hour shifts were nearly twice as likely to be identified as poor sleepers compared to those on 8-hour shifts. Finally, many more participants reported personal preference of the 8-hour over the 12-hour shift schedule. The top three issues identified as interfering with sleep were temperature, light, and noise.

**Conclusion:** Results show that sleep quality, quantity, and sleeping conditions remain problems for these military shift workers. Preliminary findings suggest that the 8-hour shift schedule is preferable to the 12-hour one, both in terms of personal preference and performance. Efforts are underway in this population to assess the use of High Energy Visible (HEV) blue light-blocking glasses to facilitate circadian entrainment and improve sleep during daytime hours. The views expressed are those of the authors and do not necessarily reflect the official policy or position of the DoN, DoD, or the U.S. Government.

**Support (If Any):**

## 0172

### PREVALENCE OF INSOMNIA AND EXCESSIVE DAYTIME SLEEPINESS IN US NAVY SAILORS

*Matsangas P, Shattuck NL*

Naval Postgraduate School, Monterey, CA

**Introduction:** The most common sleep-related complaint in civilian populations is insomnia. Workers in the U.S. Navy regularly experience significant levels of sleep deprivation and circadian misalignment due to long workdays and chronic shiftwork. This study assessed the prevalence of insomnia and elevated daytime sleepiness of crewmembers underway on a United States Navy ship while working.

**Methods:** Crewmembers (N=166, n=90 working on a fixed watchstanding schedule, n=76 on a rotating schedule) from the Reactor Department of the USS NIMITZ volunteered in this study. Sleep was assessed with actigraphy. Insomnia was assessed with the Insomnia Severity Index (ISI), and the Epworth Sleepiness Scale (ESS) was used to assess sleepiness.

**Results:** Participants slept an average  $6.75 \pm 0.94$  hours/day. ESS scores were negatively correlated with sleep duration ( $r = -0.175$ ,  $p = 0.033$ ); ISI scores were positively correlated with the number of sleep episodes per day ( $r = 0.221$ ,  $p = 0.007$ ). ESS scores (mean =  $9.91 \pm 4.66$ ) indicated that 45% of the participants had excessive daytime sleepiness—EDS (ESS score > 10). ISI scores (mean =  $11.5 \pm 5.20$ ) indicated that 66% of the participants have symptoms of insomnia (ISI score  $\geq 10$ ). Approximately 36% of the participants had EDS and comorbid insomnia symptoms, 30% had insomnia without EDS, while 9% had EDS without insomnia symptoms. The prevalence of EDS and insomnia was modulated by the type of the watchstanding schedule. For participants on the fixed schedule, the prevalence of insomnia was 57% for the EDS group and 35% for the normal sleepiness group ( $p < 0.001$ ). In the rotating schedule, however, the prevalence of insomnia was ~86% for both EDS and normal sleepiness groups ( $p > 0.9$ ).

**Conclusion:** Both excessive daytime sleepiness and insomnia remain problems for crewmembers working at sea. Results show that, as

expected, sleepiness increased with level of sleep deprivation. The prevalence of insomnia increased dramatically when crewmembers worked rotating, non-circadian schedules in which their sleep was split into multiple episodes during the day. The latter result suggests that these rotating non-circadian schedules induce insomnia symptoms due to circadian misalignment and irregular sleep patterns.

**Support (If Any):**

## 0173

### CHANGES IN SLEEPINESS RATINGS ACROSS SHIFTS AMONG MOTORCOACH DRIVERS

*Wu LJ<sup>1</sup>, Belenky G<sup>2</sup>*

<sup>1</sup>Massey University Sleep/Wake Research Centre, Wellington, NEW ZEALAND, <sup>2</sup>Washington State University Sleep and Performance Research Center, Spokane, WA

**Introduction:** The timing of shift start affects motorcoach drivers' ratings of sleepiness. Here we compare the start of work sleepiness ratings to those made at the end of work among commercial motorcoach drivers.

**Methods:** Seventy-eight commercial motorcoach drivers were monitored for approximately one month as they completed their usual work and rest schedules. Drivers kept a sleep/work diary, continuously wore an actigraph to record sleep/wake, and self-rated their sleepiness with the Karolinska Sleepiness Scale at the start and end of work periods. Sleep duration within each 24-hour period preceding duty start was summed, and shift end times were binned into morning (06:00 to 13:59), afternoon (14:00 to 21:59), and night (22:00 to 05:59). Changes in sleepiness ratings were analysed using linear mixed-effects models.

**Results:** During the study period 1,518 work periods were observed, though pre- and post-work ratings for both measures were available for only 1306 shifts. Work periods tended to end in the afternoon (mean =  $17:24 \pm 5:46$ ) and averaged  $9.2 (\pm 3.0)$  hours in duration. Drivers obtained a mean of  $6.4 (\pm 1.6)$  hours of sleep during the 24 hours prior to duty start. Subjective sleepiness ratings were highest following shifts that ended in the night. Post-shift sleepiness ratings were predicted by rating at shift start, duration of shift, and timing of shift end. Total sleep time in the 24 hours of shift start did not predict end-of-shift sleepiness ratings.

**Conclusion:** Duration of pre-work sleep did not predict sleepiness scores at the end of the shift, while operational factors such as shift length, shift timing, and the pre-work sleepiness ratings were important in determining end-of-shift sleepiness levels. Though shift timing is usually dictated by commercial demands in motorcoach operations, these data suggest that shift timing is an important consideration in managing sleepiness.

**Support (If Any):** Funded by the Federal Motor Carrier Safety Administration.

## 0174

### THERE AND BACK AGAIN: CIRCADIAN MODULATION OF SLEEP IN PILOTS FLYING ULTRA-LONG RANGE FLIGHTS

*Lamp A, Burduli E, Chen JM, Belenky G*

Washington State University, Spokane, WA

**Introduction:** Ultra Long Range (ULR; 16+ hour) flights cross multiple time zones. Therefore during layover (26–40 hours), the light/dark cycle is radically out of phase relative to home base time. The question is to what degree do pilots shift or readjust their sleep/wake cycle during layover and post-flight days as measured by synchronization to home base time.

**Methods:** Pilots' sleep/wake history was recorded by actigraph and a sleep/work logbook from three days prior to the outbound flight through three days following the inbound flight. The sleep/wake